

**DNAX RESEARCH INSTITUTE**

901 California Avenue  
Palo Alto, CA 94304-1104  
(415) 852-9196

**ASSISTANT COMMISSIONER FOR PATENTS**  
Box: Patent Application  
Washington, D. C. 20231



Transmitted herewith for filing is the:

☒ patent application of:

**MAMMALIAN CHEMOKINE REAGENTS**

Inventors: Wei Wang  
Kurt C. Gish  
Thomas J. Schall  
Alain Vicari  
Albert Zlotnik

**PATENT**

"Express Mail" Label No.: **EM 161 836 205 US**

Date of Deposit: **July 3, 1997**

Atty. Docket No.: **DX0589Q**

I hereby certify that this is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to the Commissioner of Patents and Trademarks, Box: Patent Application, Washington, D. C. 20231.

By *Lois E. Miller* *July 3, 1997*  
Lois E. Miller July 3, 1997

Sir:

Enclosed are:

☒ RETURN RECEIPT POSTCARD

☒ PATENT APPLICATION TRANSMITTAL **EM 161 836 205 US**: 1 Page (in duplicate)

☒ DECLARATION BY INVENTORS/POWER OF ATTORNEY: 3 Pages: ☐ Signed ☒ Unsigned

☒ PATENT APPLICATION TITLE PAGE: 1 Page (unnumbered)

☒ SPECIFICATION: 106 Pages: 1 through 106

☒ SEQUENCE LISTING: 22 Pages, page 107 through 128

☒ CLAIMS (22): 6 Pages: 129 through 134

☒ 1 sheet of ☐ formal ☒ informal drawing

☒ ABSTRACT: 1 Page: Page 135

☐ Appendix

☐ An assignment of the invention to Schering Corporation, a New Jersey Corporation

☐ Sequence Transmittal (in duplicate), diskette, and paper print out

**In view of the Unsigned Declaration as filed with this application and pursuant to 37 CFR § 1.53(d), Applicant requests deferral of the filing fee until submission of the Missing Parts of Application.**

Please charge DNAX Deposit Account No. 04-1239 as follows:

☒ Any additional fees associated with this paper or during the pendency of this application

-two- copies of this sheet are enclosed.

Respectfully submitted,  
DNAX RESEARCH INSTITUTE

Telephone:  
(415) 852-9196

*Edwin P. Ching* *July 3, 1997*  
Edwin P. Ching date  
Reg. No. 34,090

**PATENT APPLICATION**

5

**MAMMALIAN CHEMOKINE REAGENTS**

**INVENTORS:**

10

Wei Wang, a citizen of the Peoples' Republic of China,  
residing at 3090 Middlefield Road; Palo Alto,  
California 94306;

15

Kurt C. Gish, a citizen of the United States, residing  
at 974 Hayes Street; San Francisco, California  
94117;

20

Thomas J. Schall, a citizen of the United States,  
residing at 2050 Mills Avenue; Menlo Park,  
California 94025;

25

Alain P. Vicari, a citizen of France, residing at 2250  
Latham Street #54; Mountain View, California  
94040; and

30

Albert Zlotnik, a citizen of the United States,  
residing at 507 Alger Drive; Palo Alto,  
California 94306.

**EM161836205US**

35

Express Mail label number EM 161 836 205 US  
Date of Deposit is July 3, 1997  
I hereby certify that this paper is being deposited with the United  
States Postal Service "Express Mail Post Office to Addressee" service  
under 37 C.F.R. § 1.10 on the date indicated above and is addressed to:  
Box Patent Application; Honorable Commissioner for Patents and  
Trademarks; Washington, D.C. 20231.

40

*Lois E. Miller* *July 3, 1997*  
Lois E. Miller date

45

**Assignee:**

Schering Corporation, a New Jersey corporation

50

DNAX Research Institute  
901 California Avenue  
Palo Alto, California 94304-1104  
Tel: (415)852-9196  
Fax: (415)496-1200

## MAMMALIAN CHEMOKINE REAGENTS

This filing is a regular U.S. Patent Application resulting from the conversion of provisional applications.

## 5 FIELD OF THE INVENTION

The present invention relates to compositions related to proteins which function in controlling physiology, development, and/or differentiation of mammalian cells, e.g., cells of a mammalian immune system. In particular,  
10 it provides proteins and mimetics which regulate physiology, development, differentiation, and function of various cell types, including hematopoietic cells. It also provides receptor reagents for chemokine-like proteins.

## 15 BACKGROUND OF THE INVENTION

The circulating component of the mammalian circulatory system comprises various cell types, including red and white blood cells of the erythroid or the myeloid cell lineages. See, e.g., Rapaport (1987) Introduction to  
20 Hematology (2d ed.) Lippincott, Philadelphia, PA; Jandl (1987) Blood: Textbook of Hematology, Little, Brown and Co., Boston, MA.; and Paul (ed.) (1993) Fundamental Immunology 3d ed, Raven Press, N.Y. Progression through various stages of differentiation are regulated by various  
25 signals provided to the cells, often mediated through a class of proteins known as the cytokines. Within this group of molecules as a further group known as the chemoattractant cytokines, or chemokines. See, e.g., Schall (1994) "The Chemokines" in Thomson (ed.) The  
30 Cytokine Handbook (2d ed.) Academic Press; and Schall and Bacon (1994) Current Opinion in Immunology 6:865-873.

Although the full spectrum of biological activities of the chemokines has not been extensively investigated, chemoattractant effects are recognized. The best known  
35 biological functions of these molecules relate to chemoattraction of leukocytes. However, new chemokines are being discovered, and their biological effects on the

various cells responsible for immunological responses are topics of continued study.

Certain G-protein coupled receptors have also been characterized, presumably chemokine receptors. See, e.g.,  
5 Samson, et al. (1996) Biochemistry 35:3362-3367; and  
Rapport, et al. (1996) J. Leukocyte Biology 59:18-23.

These observations indicate that other factors exist whose functions in hematopoiesis, immune development, and leukocyte trafficking were heretofore unrecognized. These  
10 factors provide for biological activities whose spectra of effects are distinct from known differentiation, activation, or other signaling factors. The absence of knowledge about the structural, biological, and  
physiological properties of the regulatory factors which  
15 regulate hematopoietic cell physiology in vivo prevents the modification of the effects of such factors. Thus, medical conditions where regulation of the development or physiology of relevant cells is required remains unmanageable.

## SUMMARY OF THE INVENTION

The present invention is based, in part, upon the discovery of new genes encoding chemokines, and new genes encoding various receptors for chemokines. It embraces agonists and antagonists of the chemokines. In particular, sequences of various chemokines, e.g., designated Thymus Expressed ChemoKine (TECK); MIP-3 $\alpha$ ; MIP-3 $\beta$ ; and 7 transmembrane receptors, designated "dendritic cell receptor for chemokine" (DC CR) and "monocyte/dendritic cell receptor for chemokine" (M/DC CR); and mutations (muteins) of the respective natural sequences, fusion proteins, chemical mimetics, antibodies, and other structural or functional analogs are provided. It is also directed to isolated genes encoding respective proteins of the invention. Various uses of these different protein or nucleic acid compositions are also provided.

The present invention provides a substantially pure or isolated polypeptide comprising a segment exhibiting sequence homology to a corresponding portion of a mature TECK, MIP-3 $\alpha$ , MIP-3 $\beta$ , DC CR, or M/DC CR, wherein the homology is at least about 70% identity and the portion is at least about 25 amino acids. Preferably, the protein further comprises a second segment exhibiting at least about 90% identity over at least 9 amino acids; or at least about 80% identity over at least 17 amino acids. In other preferred embodiments, the polypeptide: is from a warm blooded animal selected from the group of birds and mammals, including a mouse or human; comprises a natural sequence from Tables 1 through 5; exhibits a post-translational modification pattern distinct from a natural form of the polypeptide; is made by expression of a recombinant nucleic acid; comprises synthetic sequence; is detectably labeled; is conjugated to a solid substrate; is conjugated to another chemical moiety; is a fusion protein; is in a denatured conformation, including detergent denaturation; further comprises an epitope tag; is an immunogenic polypeptide; has a defined homogeneous

molecular weight; is useful as a carbon source; is an allelic variant of SEQ ID NO: 2, 4, 6, 8, 10, or 12; is a 3-fold or less substituted form of a natural sequence; is in a sterile composition; is in a buffered solution or  
5 suspension; is in a regulated release device; comprises a post-translational modification; is in a cell; or is in a kit which further comprises instructions for use or disposal of reagents therein.

In other aspects, the invention provides an isolated  
10 or recombinant nucleic acid encoding such protein, where the portion consists of sequence from the coding region of SEQ ID NO: 1, 3, 5, 7, 9, or 11. Other aspects include such nucleic acids which: exhibit at least about 80% identity to a natural cDNA encoding said segment; is in an  
15 expression vector; further comprises a promoter; further comprises an origin of replication; is from a natural source; is detectably labeled; comprises synthetic nucleotide sequence; is less than 6 kb; is from a mammal; comprises a natural full length mature coding sequence; is  
20 in a kit, which also comprises instructions for use or disposal of reagents therein; is a specific hybridization probe for a gene encoding the protein; is a PCR product; or is in a cell. The invention also provides a method of using a purified nucleic acid by expressing the nucleic  
25 acid to produce a protein.

Alternatively, the invention provides an isolated or recombinant nucleic acid which encodes at least eight consecutive residues of SEQ ID NO: 2, 4, 6, 8, 10, or 12. Preferably, that nucleic acid encodes at least: twelve  
30 consecutive residues from SEQ ID NO: 2, and further comprises a coding sequence of at least 17 nucleotides from SEQ ID NO: 1; twelve consecutive residues from SEQ ID NO: 4, and further comprises a coding sequence of at least 17 nucleotides from SEQ ID NO: 3; twelve consecutive residues  
35 from SEQ ID NO: 6, and further comprises a coding sequence of at least 17 nucleotides from SEQ ID NO: 5; twelve consecutive residues from SEQ ID NO: 8, and further

comprises a coding sequence of at least 17 nucleotides from SEQ ID NO: 7; twelve consecutive residues from SEQ ID NO: 10, and further comprises a coding sequence of at least 17 nucleotides from SEQ ID NO: 9; or twelve consecutive  
5 residues from SEQ ID NO: 12, and further comprises a coding sequence of at least 17 nucleotides from SEQ ID NO: 11. In other preferred embodiments, the nucleic acid: exhibits at least about 80% identity to a natural cDNA encoding the segment; is in an expression vector; further comprises a  
10 promoter; further comprises an origin of replication; encodes a 3-fold or less substituted sequence from a natural sequence; is from a natural source; is detectably labeled; comprises synthetic nucleotide sequence; is less than 6 kb; is from a mammal; is attached to a solid  
15 substrate, including in a Southern or Northern blot; comprises a natural full length coding sequence; is in a cell; or is in a detection kit, which also comprises instructions for use or disposal of reagents therein. Further embodiments include a nucleic acid which hybridizes  
20 under stringent wash conditions of 55° C and less than 150 mM salt to the nucleic acid; while preferred embodiments include those which exhibit at least about 85% identity over a stretch of at least about 30 nucleotides to a primate sequence of SEQ ID NO: 1, 3, 5, 7, 9, or 11; or  
25 where the identity is at least 90%; or the stretch is at least 75 nucleotides; or where the identity is at least 95%; or the stretch is at least 100 nucleotides.

In other embodiments, the invention provides a binding compound comprising an antigen binding fragment from an  
30 antibody which binds to a mature TECK, MIP-3 $\alpha$ , MIP-3 $\beta$ , DC CR, or M/DC CR protein. In various embodiments, the binding compound is one wherein: the polypeptide is a mouse or human protein; the antibody is raised against a mature peptide sequence of Tables 1 through 5; the antibody is a  
35 monoclonal antibody; the binding compound is attached to a solid substrate; the binding compound is in a sterile composition; the binding compound binds to a denatured

antigen, including a detergent denatured antigen; the binding compound is detectably labeled; the binding compound is an Fv, Fab, or Fab2 fragment; the binding compound is conjugated to a chemical moiety; the binding  
5 compound is in a detection kit which also comprises instructions for use or disposal of reagents therein.

The invention also provides a cell which makes the antibody.

The invention embraces methods of purifying a  
10 polypeptide using a binding compound to specifically separate said polypeptides from others; of generating an antigen-binding compound complex comprising the step of contacting a sample comprising the antigen to a sample comprising a binding compound; or of modulating physiology  
15 or development of a cell expressing a receptor for a chemokine selected from TECK, MIP-3 $\alpha$ , or MIP-3 $\beta$ ; the method comprising contacting the cell with a composition comprising an agonist or mutein of said chemokine or an antibody antagonist of the chemokine. In certain  
20 embodiments of the method, the cell is a macrophage, lymphocyte, or eosinophil; or the physiology is a cellular calcium flux, a chemoattractant response, cellular morphology modification responses, phosphoinositide lipid turnover, or an antiviral response. In other embodiments,  
25 the receptor is DC CR and the chemokine is MIP-3 $\alpha$ , the physiology is pulmonary physiology, or the cell is an eosinophil.



## DESCRIPTION OF THE DRAWINGS

Figures 1A-1B show chemotactic properties of mTECK recombinant protein. Fig. 1A shows migration of mouse thymocytes to recombinant mTECK and effect of pertussis toxin. Chemotaxis assays were performed as described. Recombinant mouse lymphotactin was used as a positive control. Data are expressed as the mean of cell counts obtained from three separate experiments in duplicate  $\pm$  SEM. In one experiment, cells were pre incubated 1 h with 10 ng/ml pertussis toxin (PTX) prior to the assay. Fig. 1B shows migration of other leukocyte subsets to recombinant mTECK. Mouse splenic dendritic cells and mouse activated macrophages were obtained. THP-1 human monocytic cells were used without or with a 16 h activation with IFN- $\gamma$ . Results are obtained as the mean of the chemotactic index from three separate experiments per cell type in duplicate  $\pm$  SD. The number of cells migrating to medium alone was greater than 40 cells per 5 high power fields in each experiment. Recombinant MIP-1 $\alpha$  was used as a positive control.

## DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

## OUTLINE

- 5
- I. General
  - II. Purified Chemokines, Receptors
    - A. physical properties
    - B. biological properties
  - 10 III. Physical Variants
    - A. sequence variants, fragments
    - B. post-translational variants
      - 1. glycosylation
      - 2. others
  - 15 IV. Functional Variants
    - A. analogs; fragments
      - 1. agonists
      - 2. antagonists
    - B. mimetics
      - 1. protein
      - 2. chemicals
    - C. species variants
  - 20 V. Antibodies
    - A. polyclonal
    - 25 B. monoclonal
    - C. fragments, binding compositions
  - VI. Nucleic Acids
    - A. natural isolates; methods
    - B. synthetic genes
    - 30 C. methods to isolate
  - VII. Making Chemokines, Receptors; Mimetics
    - A. recombinant methods
    - B. synthetic methods
    - C. natural purification
  - 35 VIII. Uses
    - A. diagnostic
    - B. therapeutic
  - IX. Kits
    - A. nucleic acid reagents
    - 40 B. protein reagents
    - C. antibody reagents
  - X. Receptors

## I. General

45 The present invention provides DNA sequences encoding various mammalian proteins which exhibit structural properties characteristic of a chemotactic cytokine, or chemokine. Other embodiments are directed to chemokine receptors. See, e.g., Lodi, et al. (1994) Science  
50 263:1762-1767; Gronenborn and Clore (1991) Protein

- Engineering 4:263-269; Miller and Kranger (1992) Proc. Nat'l Acad. Sci. USA 89:2950-2954; Matsushima and Oppenheim (1989) Cytokine 1:2-13; Stoeckle and Baker (1990) New Biol. 2:313-323; Oppenheim, et al. (1991) Ann. Rev. Immunol. 9:617-648; Schall (1991) Cytokine 3:165-183; and The Cytokine Handbook Academic Press, NY. Mouse and human embodiments are described herein.

Chemokines play an important role in immune and inflammatory responses by inducing migration and adhesion of leukocytes. These small secreted molecules are a growing superfamily of 8-14 kDa proteins characterized by a conserved four cysteine motif. See, e.g., Schall (1991) Cytokine 3:165-183; and Thomson (ed.) The Cytokine Handbook Academic Press, NY. Chemokines are secreted by activated leukocytes and act as a chemoattractant for a variety of cells which are involved in inflammation. Besides chemoattractant properties, chemokines have been shown to induce other biological responses, e.g., modulation of second messenger levels such as  $Ca^{++}$ ; inositol phosphate pool changes (see, e.g., Berridge (1993) Nature 361:315-325 or Billah and Anthes (1990) Biochem. J. 269:281-291); cellular morphology modification responses; phosphoinositide lipid turnover; possible antiviral responses; and others. Thus, the chemokines provided herein may, alone or in combination with other therapeutic reagents, have advantageous combination effects.

Moreover, there are reasons to suggest that chemokines may have effects on other cell types, e.g., attraction or activation of monocytes, dendritic cells, T cells, eosinophils, and/or perhaps on basophils and/or neutrophils. They may also have chemoattractive effects on various neural cells including, e.g., dorsal root ganglia neurons in the peripheral nervous system and/or central nervous system neurons.

Membrane proteins which contain seven transmembrane segments have been characterized as G-protein coupled receptors. Many of these receptors have been characterized

as receptors for chemokines, based in part on structural features. Chemokine receptors are important in the signal transduction mechanisms mediated by the chemokines. They are useful markers for distinguishing cell populations, and  
5 have been implicated as specific receptors for retroviral infections.

The chemokine superfamily was classically divided into two groups exhibiting characteristic structural motifs, the Cys-X-Cys (C-X-C) and Cys-Cys (C-C) families. These were  
10 distinguished on the basis of a single amino acid insertion between the NH-proximal pair of cysteine residues and sequence similarity. Typically, the C-X-C chemokines, i.e., IL-8 and MGSA/Gro- $\alpha$  act on neutrophils but not on monocytes, whereas the C-C chemokines, i.e., MIP-1 $\alpha$  and  
15 RANTES, are potent chemoattractants for monocytes and lymphocytes but not neutrophils. See, e.g., Miller, et al. (1992) Crit. Rev. Immunol. 12:17-46. A recently isolated chemokine, lymphotactin, does not belong to either group and may constitute a first member of a third chemokine  
20 family, the C family. Lymphotactin does not have a characteristic CC or CXC motif, and acts on lymphocytes but not neutrophils and monocytes. See, e.g., Kelner et al. (1994) Science 266:1395-1399. This chemokine defines a new C-C chemokine family. Even more recently, another  
25 chemokine exhibiting a CX3C motif has been identified, which establishes a fourth structural class.

The present invention provides additional chemokine reagents, e.g., nucleic acids, proteins and peptides, antibodies, etc., related to the newly discovered  
30 respective chemokines designated TECK; MIP-3 $\alpha$ , and MIP-3 $\beta$ .

In other embodiments, the invention provides two genes encoding novel 7-transmembrane (7-TM) receptors, presumably G-protein coupled receptors and likely chemokine receptors. These 7-TM receptors are hypothesized to be chemokine  
35 receptors and have been designated DC CR and M/DC CR. Their ligands have not yet specifically been completely identified. However, the receptors exhibit structural

features typical of known chemokine receptors, e.g., 7  
transmembrane spanning structures. They may exhibit  
properties of binding many different cytokines at varying  
specificities (shared or promiscuous binding specificity)  
5 or may exhibit high affinity for one (specific) or a subset  
(shared) of chemokines.

The described chemokines and receptors should be  
important for mediating various aspects of cellular, organ,  
tissue, or organismal physiology or development.

10

## II. Purified chemokines, receptors

Mouse and human Thymus Expressed ChemoKine (TECK)  
nucleotide and amino acid sequences are shown in Table 1.  
Nucleotide and amino acid sequences of another novel  
15 chemokine, from human, designated MIP-3 $\alpha$  are provided in  
Table 2. Nucleotide and derived amino acid sequences of a  
third novel chemokine, from human, designated MIP-3 $\beta$  are  
shown in Table 3. Generic descriptions of physical  
properties of polypeptides, nucleic acids, and antibodies  
20 where directed to one embodiment clearly are generally  
applicable to other chemokines or receptors described  
herein.

The nucleotide and amino acid sequences of a novel  
chemokine receptor found on dendritic cells (DC), from  
25 human, and designated DC CR, are provided in Table 4. The  
nucleotide and amino acid sequences of another novel  
chemokine receptor found on macrophages and dendritic  
cells, from human, and designated M/DC CR, are provided in  
Table 5.

30 These amino acid sequences, provided amino to carboxy,  
are important in providing sequence information on the  
chemokine ligand or receptor, allowing for distinguishing  
the protein from other proteins. Moreover, the sequences  
allow preparation of peptides to generate antibodies to  
35 recognize and distinguish such segments, and allow  
preparation of oligonucleotide probes, both of which are  
strategies for isolation, e.g., cloning, of genes encoding

- such sequences, or related sequences, e.g., natural polymorphic or other variants. Similarities of the chemokines have been observed with other cytokines. See, e.g., Bosenberg, et al. (1992) Cell 71:1157-1165; Huang, et. al. (1992) Molecular Biology of the Cell 3:349-362; and Pandiella, et al. (1992) J. Biol. Chem. 267:24028-24033.
- 5 Likewise for the receptors.

Table 1: Nucleotide sequence (5' to 3') of TECK from mouse and the corresponding amino acid sequence (amino to carboxy). Signal sequence probably runs as shown between Ala and Gln, see SEQ ID NO: 1 and 2. Human sequences are SEQ ID NO: 3 and 4.

5	1	AGGCTACAAGCAGGCACCAGCTCTCAGGACCAGAAAGGCATTGGTGGCCCCCTTAAACCT	60
10	61	TCAGGTATCTGGAGAGGAGATCTAACCTTCACTATGAAACTGTGGCTTTTGCCTGCCTG	120
	1	MetLysLeuTrpLeuPheAlaCysLeu	9
	121	GTTGCCTGTTTTGTTGGGGCCTGGATGCCGGTTGTCCATGCCCAAGGTGCCTTTGAAGAC	180
	10	ValAlaCysPheValGlyAlaTrpMetProValValHisAlaGlnGlyAlaPheGluAsp	29
15	181	TGCTGCCTGGGTTACCAGCACAGGATCAAATGGAATGTTCTCCGGCATGCTAGGAATTAT	240
	30	CysCysLeuGlyTyrGlnHisArgIleLysTrpAsnValLeuArgHisAlaArgAsnTyr	49
	241	CACCAGCAGGAAGTGAGTGAAGCTGCAACCTACGTGCTGTGAGATTCTACTTCCGCCAG	300
20	50	HisGlnGlnGluValSerGlySerCysAsnLeuArgAlaValArgPheTyrPheArgGln	69
	301	AAAGTAGTGTGTGGGAATCCAGAGGACATGAATGTGAAGAGGGCGATAAGAATCTTGACA	360
	70	LysValValCysGlyAsnProGluAspMetAsnValLysArgAlaIleArgIleLeuThr	89
25	361	GCTAGGAAAAGGCTAGTCCACTGGAAGAGCGCCTCAGACTCTCAGACTGAAAGGAAGAAG	420
	90	AlaArgLysArgLeuValHisTrpLysSerAlaSerAspSerGlnThrGluArgLysLys	109
	421	TCAAACCATATGAAGTCCAAGGTGGAGAACCCCAACAGTACAAGCGTGAGGAGTGCCACC	480
	110	SerAsnHisMetLysSerLysValGluAsnProAsnSerThrSerValArgSerAlaThr	129
30	481	CTAGGTCATCCCAGGATGGTGATGATGCCCAGAAAGACCAACAATTAAGTTAATTACTCA	540
	130	LeuGlyHisProArgMetValMetMetProArgLysThrAsnAsnEnd	144
	541	GAGTAAGCACCAGCTGGAGGATGGGCGGAGTCTGCTGAAGTGCTGTCTTCTAGGCATGCC	600
35	601	AGTGCCAATGAACTCACTGAAGCTACAGTTTCCTGTACAAGACCAGACCCACCAACGTCT	660
	661	CAGCATGTACGAGGAAGGAAGTACTGCGCTAAAGGCCCTCCCACTCACCAAGGAGCTATT	720
40	721	GGCTATTGATGATTGCTGAGGGAAGGGAGTAATTTTTTTTCTCTTTCTGAAGTGTGACTT	780
	781	GAGTAAATTGCCCATAGTTTCAGTATATAATCCCCAACCTGTGCTCAGGCAAGCAACCCTA	840
	841	ATTAAATGCAATAGCCACATACAAAAGAAGAGGATATGAATAGTTTGGTAGGAGGGGCTT	900
45	901	GTTAGGAAGAAGACATTAACAGGAGAGAGAGGAGCGAGAGGATAGTGAGTGTGTGAGAGT	960
	961	GCCTGCACGTGTGAAATGGTCAAAGAATAAAAAATAAAAACTTAAAAAGCTATTAAAAA	1020
50	1021	GTAAAAAAAATAAA 1034	

Table 1 (continued):

human Teck cDNA (see SEQ ID NO: 3); signal sequence cleavage is probably between about Thr and Gln. Hu TECK protein sequence (see SEQ ID NO: 4).

5	TCGACCCACG CGTCCGCTTG GCCTACAGCC CGGCGGGCAT CAGCTCCCTT GACCCAGTGG	60
	ATATCGGTGG CCCC GTTATT CGTCCAGGTG CCCAGGGAGG AGGACCCGCC TGCAGC	116
10	ATG AAC CTG TGG CTC CTG GCC TGC CTG GTG GCC GGC TTC CTG GGA GCC Met Asn Leu Trp Leu Leu Ala Cys Leu Val Ala Gly Phe Leu Gly Ala -23 -20 -15 -10	164
15	TGG GCC CCC GCT GTC CAC ACC CAA GGT GTC TTT GAG GAC TGC TGC CTG Trp Ala Pro Ala Val His Thr Gln Gly Val Phe Glu Asp Cys Cys Leu -5 1 5	212
20	GCC TAC CAC TAC CCC ATT GGG TGG GCT GTG CTC CGG CGC GCC TGG ACT Ala Tyr His Tyr Pro Ile Gly Trp Ala Val Leu Arg Arg Ala Trp Thr 10 15 20 25	260
25	TAC CGG ATC CAG GAG GTG AGC GGG AGC TGC AAT CTG CCT GCT GCG ATA Tyr Arg Ile Gln Glu Val Ser Gly Ser Cys Asn Leu Pro Ala Ala Ile 30 35 40	308
30	TTC TAC CTC CCC AAG AGA CAC AGG AAG GTG TGT GGG AAC CCC AAA AGC Phe Tyr Leu Pro Lys Arg His Arg Lys Val Cys Gly Asn Pro Lys Ser 45 50 55	356
35	AGG GAG GTG CAG AGA GCC ATG AAG CTC CTG GAT GCT CGA AAT AAG GTT Arg Glu Val Gln Arg Ala Met Lys Leu Leu Asp Ala Arg Asn Lys Val 60 65 70	404
40	TTT GCA AAG CTC CAC CAC AAC ATG CAG ACC TTC CAA GCA GGC CCT CAT Phe Ala Lys Leu His His Asn Met Gln Thr Phe Gln Ala Gly Pro His 75 80 85	452
45	GCT GTA AAG AAG TTG AGT TCT GGA AAC TCC AAG TTA TCA TCA TCC AAG Ala Val Lys Lys Leu Ser Ser Gly Asn Ser Lys Leu Ser Ser Ser Lys 90 95 100 105	500
50	TTT AGC AAT CCC ATC AGC AGC AGC AAG AGG AAT GTC TCC CTC CTG ATA Phe Ser Asn Pro Ile Ser Ser Ser Lys Arg Asn Val Ser Leu Leu Ile 110 115 120	548
55	TCA GCT AAT TCA GGA CTG TGAGCCGGCT CATTTCTGGG CTCCATCGGC Ser Ala Asn Ser Gly Leu 125	596
	ACAGGAGGGG CCGGATCTTT CTCCGATAAA ACCGTGCCCC TACAGACCCA GCTGTCCCCA	656
	CGCCTCTGTC TTTTGGGTCA AGTCTTAATC CCTGCACCTG AGTTGGTCCT CCCTCTGCAC	716
	CCCCACCACC TCCTGCCCCG CTGGCAACTG GAAAGAAGGA GTTGGCCTGA TTTTAACCTT	776
	TTGCCGCTCC GGGGAACAGC ACAATCCTGG GCAGCCAGTG GCTCTTGTAG AGAAACTTA	836
	GGATACCTCT CTCACCTTCT GTTTCTTGCC GTCCACCCCG GGCCATGCCA GTGTGTCTCTC	896



Table 1 (continued):

	TGGGTCCCCT CCAAAAATCT GGTCAATCAA GGATCCCCTC CCAAGGCTAT GCTTTTCTAT	956
5	AACTTTTAAA TAAACCTTGG GGGGTGAATG GAATAAAAAA AAAAAAAAAA AAAAAA	1012

10 Table 2: Nucleotide sequence (5' to 3') of MIP-3 $\alpha$  from human and the corresponding amino acid sequence (amino to carboxy), see SEQ ID NO: 5 and 6 and GenBank Accession U77035.

	ATG TGC TGT ACC AAG AGT TTG CTC CTG GCT GCT TTG ATG TCA GTG CTG	48
15	Met Cys Cys Thr Lys Ser Leu Leu Ala Ala Leu Met Ser Val Leu -26 -25 -20 -15	
	CTA CTC CAC CTC TGC GGC GAA TCA GAA GCA GCA AGC AAC TTT GAC TGC	96
20	Leu Leu His Leu Cys Gly Glu Ser Glu Ala Ala Ser Asn Phe Asp Cys -10 -5 1 5	
	TGT CTT GGA TAC ACA GAC CGT ATT CTT CAT CCT AAA TTT ATT GTG GGC	144
	Cys Leu Gly Tyr Thr Asp Arg Ile Leu His Pro Lys Phe Ile Val Gly 10 15 20	
25	TTC ACA CGG CAG CTG GCC AAT GAA GGC TGT GAC ATC AAT GCT ATC ATC	192
	Phe Thr Arg Gln Leu Ala Asn Glu Gly Cys Asp Ile Asn Ala Ile Ile 25 30 35	
30	TTT CAC ACA AAG AAA AAG TTG TCT GTG TGC GCA AAT CCA AAA CAG ACT	240
	Phe His Thr Lys Lys Lys Leu Ser Val Cys Ala Asn Pro Lys Gln Thr 40 45 50	
35	TGG GTG AAA TAT ATT GTG CGT CTC CTC AGT AAA AAA GTC AAG AAC ATG	288
	Trp Val Lys Tyr Ile Val Arg Leu Leu Ser Lys Lys Val Lys Asn Met 55 60 65 70	
	TAAAACTGT GGCTTTTCTG GAATGGAATT GGACATAGCC CAAGAACAGA AAGAACCTTG	348
40	CTGGGGTTGG AGGTTTCACT TGCACATCAT GGAGGGTTTA GTGCTTATCT AATTTGTGCC	408
	TCACTGGACT TGTCCAATTA ATGAAGTTGA TTCATATTGC ATCATAGTTT GCTTTGTTTA	468
	AGCATCACAT TAAAGTTAAA CTGTATTTTA TGTTATTTAT AGCTGTAGGT TTTCTGTGTT	528
45	TAGCTATTTA ATACTAATTT TCCATAAGCT ATTTTGGTTT AGTGCAAAGT ATAAAATTAT	588
	ATTTGGGGGG GAATAAGATT ATATGGACTT TTTTGCAAGC AACAAGCTAT TTTTAAAAA	648
50	AACTATTTA ACATTCTTTT GTTTATATTG TTTTGTCTCC TAAATTGTTG TAATTGCATT	708
	ATAAAATAAG AAAAATATTA ATAAGACAAA TATTGAAAT AAAGAAACAA AAAGTTAAAA	768
	AAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAA	801

Table 3: Nucleotide sequence (5' to 3') of MIP-3 $\beta$  from human and the corresponding amino acid sequence (amino to carboxy), see SEQ ID NO: 7 and 8, and GenBank Accession U77180. Signal sequence cleavage is about between Ser and Gly.

5	1	GGCACGAGCGGCACGAGCATCACTCACACCTTGCAATTCACCCCTGCATCCCAGTCGCCC	60
	61	TGCAGCCTCACACAGATCCTGCACACACCCAGACAGCTGGCGCTCACACATTCACCGTTG	120
10	121	GCCTGCCTCTGTTCACCCCTCCATGGCCCTGCTACTGGCCCTCAGCCTGCTGGTTCTCTGG	180
	1	MetAlaLeuLeuLeuAlaLeuSerLeuLeuValLeuTrp	13
	181	ACTTCCCCAGCCCCAACTCTGAGTGGCACCAATGATGCTGAAGACTGCTGCCTGTCTGTG	240
15	14	ThrSerProAlaProThrLeuSerGlyThrAsnAspAlaGluAspCysCysLeuSerVal	33
	241	ACCCAGAAACCCATCCCTGGGTACATCGTGAGGAATTCCACTACCTTCTCATCAAGGAT	300
	34	ThrGlnLysProIleProGlyTyrIleValArgAsnPheHisTyrLeuLeuIleLysAsp	53
	301	GGCTGCAGGGTGCCTGCTGTAGTGTTCACCACACTGAGGGGCCGCCAGCTCTGTGCACCC	360
20	54	GlyCysArgValProAlaValValPheThrThrLeuArgGlyArgGlnLeuCysAlaPro	73
	361	CCAGACCAGCCCTGGGTAGAACGCATCATCCAGAGACTGCAGAGGACCTCAGCCAAGATG	420
	74	ProAspGlnProTrpValGluArgIleIleGlnArgLeuGlnArgThrSerAlaLysMet	93
25	421	AAGCGCCGACAGCTTAACCTATGACCGTGCAGAGGGAGCCCGGAGTCCGAGTCAAGCAT	480
	94	LysArgArgSerSerEnd	98
	481	TGTGAATTATTACCTAACCTGGGGAACCGAGGACCAGAAGGAAGGACCAGGCTTCCAGCT	540
30	541	CCTCTGCACCAGACCTGACCAGCCAGGACAGGGCCTGGGGTGTGTGTGAGTGTGAGTGTG	600
	601	AGCGAGAGGGTGAGTGTGGTCTAGAGTAAAGCTGCTCCACCCCCAGATTGCAATGCTACC	660
35	661	AATAAAGCCGCCTGGTGTTTACAACCTAAAAAAAAAAAAA	699

2007/06/20 14:00:00

Table 4: Nucleotide sequence (5' to 3') of chemokine receptor, DC CR, from human and the corresponding amino acid sequence (amino to carboxy), see SEQ ID NO: 9 and 10. Nucleotide 579 may be A, C, G, or T, and the codon may code for His or Gln.

5	1	ATGTTTTCGACTCCAGTGAAGATTATTTTGTGTCAGTCAATACTTCATATTACTCAGTTG	60
	1	MetPheSerThrProValLysIleIleLeuCysGlnSerIleLeuHisIleThrGlnLeu	20
10	61	ATTCTGAGATGTTACTGTGCTCCTTGCAGGAGGTCAGGCAGTTCTCCAGGCTATTTGTAC	120
	21	IleLeuArgCysTyrCysAlaProCysArgArgSerGlySerSerProGlyTyrLeuTyr	40
15	121	CGAATTGCCTACTCCTTGATCTGTGTTCTTGGCCTCCTGGGGAATATTCTGGTGGTGATC	180
	41	ArgIleAlaTyrSerLeuIleCysValLeuGlyLeuLeuGlyAsnIleLeuValValIle	60
20	181	ACCTTTGCTTTTTATAAGAAGGCCAGGTCTATGACAGACGTCTATCTCTTGAACATGGCC	240
	61	ThrPheAlaPheTyrLysLysAlaArgSerMetThrAspValTyrLeuLeuAsnMetAla	80
25	241	ATTGCAGACATCCTCTTTGTTCTTACTCTCCCATTCTGGGCAGTGAGTCATGCCACTGGT	300
	81	IleAlaAspIleLeuPheValLeuThrLeuProPheTrpAlaValSerHisAlaThrGly	100
30	301	GCGTGGGTTTTTCAGCAATGCCACGTGCAAGTTGCTAAAAGGCATCTATGCCATCAACTTT	360
	101	AlaTrpValPheSerAsnAlaThrCysLysLeuLeuLysGlyIleTyrAlaIleAsnPhe	120
35	361	AACTGCGGGATGCTGCTCCTGACTTGCATTAGCATGGACCGGTACATCGCCATTGTACAG	420
	121	AsnCysGlyMetLeuLeuLeuThrCysIleSerMetAspArgTyrIleAlaIleValGln	140
40	421	GCGACTAAGTCATTCCGGCTCCGATCCAGAACACTACCGCGCAGCAAAATCATCTGCCTT	480
	141	AlaThrLysSerPheArgLeuArgSerArgThrLeuProArgSerLysIleIleCysLeu	160
45	481	GTTGTGTGGGGCTGTCACTCATCATCTCCAGCTCAACTTTTGTCTTCAACCAAAAATAC	540
	161	ValValTrpGlyLeuSerValIleIleSerSerSerThrPheValPheAsnGlnLysTyr	180
50	541	AACACCCAAGGCAGCGATGTCTGTGAACCCAAGTACCAAACTGTCTCGGAGCCCATCAGG	600
	181	AsnThrGlnGlySerAspValCysGluProLysTyrGlnThrValSerGluProIleArg	200
55	601	TGGAAGCTGCTGATGTTGGGGCTTGAGCTACTCTTTGGTTTCTTTATCCCTTTGATGTTT	660
	201	TrpLysLeuLeuMetLeuGlyLeuGluLeuLeuPheGlyPhePheIleProLeuMetPhe	220
60	661	ATGATATTTTGTACACGTTTCATTGTCAAAACCTTGGTGCAAGCTCAGAATTCATAAAGG	720
	221	MetIlePheCysTyrThrPheIleValLysThrLeuValGlnAlaGlnAsnSerLysArg	240
65	721	CACAAAGCCATCCGTGTAATCATAGCTGTGGTGTGTTTCTGGCTTGTGAGATTTCCT	780
	241	HisLysAlaIleArgValIleIleAlaValValLeuValPheLeuAlaCysGlnIlePro	260
70	781	CATAACATGGTCCTGCTTGTGACGGCTGCTAATTTGGGTAAAATGAACCGATCCTGCCAG	840
	261	HisAsnMetValLeuLeuValThrAlaAlaAsnLeuGlyLysMetAsnArgSerCysGln	280
75	841	AGCGAAAAGCTAATTGGCTATACGAAAAGTGTACAGAAAGTCCTGGCTTTCCTGCACTGC	900
	281	SerGluLysLeuIleGlyTyrThrLysThrValThrGluValLeuAlaPheLeuHisCys	300
80	901	TGCCTGAACCCTGTGCTCTACGCTTTTATTGGGCAGAACTTCAGAACTACTTTCTGAAG	960
	301	CysLeuAsnProValLeuTyrAlaPheIleGlyGlnLysPheArgAsnTyrPheLeuLys	320
85	961	ATCTTGAAGGACCTGTGGTGTGTGAGAAGGAAGTACAAGTCCTCAGGCTTCTCCTGTGCC	1020
	321	IleLeuLysAspLeuTrpCysValArgArgLysTyrLysSerSerGlyPheSerCysAla	340
90	1021	GGGAGGTACTCAGAAAACATTTCTCGGCAGACCAGTGAGACCGCAGATAACGACAATGCG	1080
	341	GlyArgTyrSerGluAsnIleSerArgGlnThrSerGluThrAlaAspAsnAspAla	360

Table 4 (continued)

1081 TCGTCCTTCACTATGTGATAGAAAGCTGAGTCTCCCTAA 1119  
 361 SerSerPheThrMetEnd 365

Table 5: Nucleotide sequence (5' to 3') of chemokine receptor, M/DC CR, from human and the corresponding amino acid sequence (amino to carboxy), see SEQ ID NO: 11 and 12.

10	1	GAGGAAGCTGCTTCGGGGGGTGAGCAAACCTTTTTTAAAATGCAGAAATTATGATCTACACC	60
		MetIleTyrThr	4
15	61	CGTTTCTTAAAAGGCAGTCTGAAGATGGCCAATTACACGCTGGCACCAGAGGATGAATAT	120
	5	ArgPheLeuLysGlySerLeuLysMetAlaAsnTyrThrLeuAlaProGluAspGluTyr	24
	121	GATGTCCTCATAGAAGGTGAAGTGGAGAGCGATGAGGCAGAGCAATGTGACAAGTATGAC	180
	25	AspValLeuIleGluGlyGluLeuGluSerAspGluAlaGluGlnCysAspLysTyrAsp	44
20	181	GCCCAGGCACTCTCAGCCCAGCTGGTGCCATCACTCTGCTCTGCTGTGTTGTGATCGGT	240
	45	AlaGlnAlaLeuSerAlaGlnLeuValProSerLeuCysSerAlaValPheValIleGly	64
	241	GTCCTGGACAATCTCCTGGTTGTGCTTATCCTGGTAAAATATAAAGGACTCAAACGCGTG	300
25	65	ValLeuAspAsnLeuLeuValValLeuIleLeuValLysTyrLysGlyLeuLysArgVal	84
	301	GAAAATATCTATCTTCTAAACTTGGCAGTTTCTAACTTGTGTTTCTTGCTTACCCTGCCC	360
	85	GluAsnIleTyrLeuLeuAsnLeuAlaValSerAsnLeuCysPheLeuLeuThrLeuPro	104
30	361	TTCTGGGCTCATGCTGGGGGCGATCCCATGTGTAAAATTCTCATTGGACTGTACTTCGTG	420
	105	PheTrpAlaHisAlaGlyGlyAspProMetCysLysIleLeuIleGlyLeuTyrPheVal	124
	421	GGCCTGTACAGTGAGACATTTTTCAATTGCCTTCTGACTGTGCAAAGGTACCTAGTGT	480
	125	GlyLeuTyrSerGluThrPhePheAsnCysLeuLeuThrValGlnArgTyrLeuValPhe	144
35	481	TTGCACAAGGGCAACTTTTTCTCAGCCAGGAGGAGGGTGCCCTGTGGCATCATTACAAGT	540
	145	LeuHisLysGlyAsnPhePheSerAlaArgArgValProCysGlyIleIleThrSer	164
	541	GTCCTGGCATGGGTAACAGCCATTCTGGCCACTTTGCCTGAATTCGTGGTTTATAAACCT	600
40	165	ValLeuAlaTrpValThrAlaIleLeuAlaThrLeuProGluPheValValTyrLysPro	184
	601	CAGATGGAAGACCAGAAATACAAGTGTGCATTTAGCAGAAGTCCCTTCCTGCCAGCTGAT	660
	185	GlnMetGluAspGlnLysTyrLysCysAlaPheSerArgThrProPheLeuProAlaAsp	204
45	661	GAGACATTCTGGAAGCATTCTTCTGACTTTAAAAATGAACATTTTCGGTTCTGTCTCCTCCC	720
	205	GluThrPheTrpLysHisPheLeuThrLeuLysMetAsnIleSerValLeuValLeuPro	224
	721	CTATTTATTTTACATTTCTCTATGTGCAAATGAGAAAAACACTAAGGTTTCAGGGAGCAG	780
	225	LeuPheIlePheThrPheLeuTyrValGlnMetArgLysThrLeuArgPheArgGluGln	244
50	781	AGGTATAGCCTTTTCAAGCTTGTTTTTGCCGTAATGGTAGTCTTCCTTCTGATGTGGGCG	840
	245	ArgTyrSerLeuPheLysLeuValPheAlaValMetValValPheLeuLeuMetTrpAla	264
	841	CCCTACAATATTGCATTTTCTCTGTCCACTTTCAAAGAACACTTCTCCCTGAGTGACTGC	900
55	265	ProTyrAsnIleAlaPhePheLeuSerThrPheLysGluHisPheSerLeuSerAspCys	284
	901	AAGAGCAGCTACAATCTGGACAAAAGTGTTACATCACTAAACTCATCGCCACCACCCAC	960
	285	LysSerSerTyrAsnLeuAspLysSerValHisIleThrLysLeuIleAlaThrThrHis	304
60	961	TGCTGCATCAACCCTCTCCTGTATGCGTTTCTTGATGGGACATTTAGCAAATACCTCTGC	1020
	305	CysCysIleAsnProLeuLeuTyrAlaPheLeuAspGlyThrPheSerLysTyrLeuCys	324

Table 5 (continued) M/DC CR sequence

1021	CGCTGTTTCCATCTGCGTAGTAACACCCCACTTCAACCCAGGGGGCAGTCTGCACAAGGC	1080
325	ArgCysPheHisLeuArgSerAsnThrProLeuGlnProArgGlyGlnSerAlaGlnGly	344
1081	ACATCGAGGGAAGAACCTGACCATTCCACCGAAGTGTAAGTAACTAGCATCCACCAAATGCAA	1140
345	ThrSerArgGluGluProAspHisSerThrGluValEnd	356
1141	GAAGAATAAACATGGATTTTTCATCTTTCTGCATTATTTTCATGTAAATTTTCTACACATTT	1200
1201	GTATACAAAATCGGATACAGGAAGAAAAGGGAGAGGTGAGCTAACATTTGCTAAGCACTG	1260
1261	AATTTGTCTCAGGCACCGTGCAAGGCTCTTTACAAACGTGAGCTCCTTCGCCTCCTACCA	1320
1321	CTTGTCATAGTGTGGATAGGACTAGTCTCATTTCTCTGAGAAGAAAATAAGGCGCGGA	1380
1381	AATTTGTCTAAGATCACATAACTAGGAAGTGGCAGAACTGATTCTCCAGCCCTGGTAGCA	1440
1441	TTTGCTCAGAGCCTACGCTTGGTCCAGAACATCAAACCTCCAAACCCTGGGGACAAACGAC	1500
1501	ATGAAATAAATGTATTTTAAACATATAAAAAAAAAAAAAAAAAAAAAA	1547

Table 6: Alignment of M/DC CR with CKR-1 through CKR-4. The other chemokine receptors are SEQ ID NO: 13-17. An asterisk indicates fully conserved residue among all five receptors; a period represents conservative substitutions among all five receptors.

30	M/DC CR	MIYTRFLKGS LKMAN YTLAP EDEYD VLI EGEL ESDEAE QCDKY DAQALS
	C-C CKR-1	METPNTTEDYDTTTEFDYGDATPCQKVNERAFG
	C-C CKR-2	MLSTSRSRFIRNTNESGEEVTTFFDYDYGAPCHKFDVKQIG
	C-C CKR-3	MTTSLDTVETFGTTSYYDDVGLLCEKADTRALM
	C-C CKR-4	MNPTDIADTTLDESIYSNYYLYESIPKPCTKEGIKAFG
35		* * . . .
	M/DC CR	AQLVPSLCSAVFVIGVLDNLLVVLILVKYKGLKRVENIYLLNLAVSNLCF
	C-C CKR-1	AQLLPPLYSLVFVIGLVGNILVVLVQYKRLKNMTSIYLLNLAISDLLF
	C-C CKR-2	AQLLPPLYSLVFIFGFVGNMLVVLILINCKKLKCLTDIYLLNLAISDLLF
40	C-C CKR-3	AQFVPPLYSLVFTVGLLGNNVVMILIKYRRLRIMTNIYLLNLAISDLLF
	C-C CKR-4	ELFLPPLYSLVFVFGLLGNSVVVLVLFKYKRLRSMTDVYLLNLAISDLLF
		..*.* * ** * ..* ..* * . . . * . . . *****.* * *
	M/DC CR	LLTLPFWAHAG-----GDPMCKILIGLYFVGLYSETFFNCLLTVQRYL
45	C-C CKR-1	LFTLPFWIDYKLKDDWVFGDAMCKILSGFYITGLYSEIFFIILLTIDRYL
	C-C CKR-2	LITLPLWAH-SANNEWVFGNAMCKLFTGLYHIGYFGGIFFIILLTIDRYL
	C-C CKR-3	LVTLPFWIHVVRGHNWVFGHGMCKLLSGFYHTGLYSEIFFIILLTIDRYL
	C-C CKR-4	VFSLPFWGYA-ADQWVFGGLCKMISWMYLVGFYSGIFFVMLMSIDRYL
		. .***.* * ..* * . . . * * . . . ** * . . . .***
50	M/DC CR	VFLHKGNNFFSAR-RRVPCGIITSVLAWVTAILATLPEFVVYKPMEDQKY
	C-C CKR-1	AIVH--AVFALRARTVTFGVITSIIWALAILASMPGLYFSKTQWEFTHH
	C-C CKR-2	AIVH--AVFALKARTVTFGVVTSVITWLAVFASVPGIIFTKCKEDSVY
	C-C CKR-3	AIVH--AVFALRARTVTFGVITSIVTWGLAVLAALPEFIFYETEELFEET
55	C-C CKR-4	AIVH--AVFSLRARTLTYGVITSLATWSVAVFASLPGLFSTCYTERNHT
		..* * . . . * . . . . * * . . . . * . . . .

Table 6 (continued)

	M/DC CR	KCAFSRTPFLPADETF-WKHFLTLKMNISVLVLPLFIFTFLYVQMRKTL-
	C-C CKR-1	TCS----LHFPHESLREWKLFQALKLNLFGLVLP LLVMIICYTGIIKILL
5	C-C CKR-2	VCG----PYFPR---GWN NFHTIMRNILGLVLP LLIMVICYSGILKTLL
	C-C CKR-3	LCS----ALYPEDTVYSWRHFHTLRMTIFCLVLP LLVMAICYTGIIKTLL
	C-C CKR-4	YCK----TKYSLNST-TWKVLSSLEINILGLVLP LGIMLFCYSMIIRTLO
		* . . . . * . . . . * . . . . *
10	M/DC CR	--RFREQRYSLFKLVFAVMVVFLLMWAPYNIAFFLSTFKEHFSLS DCKSS
	C-C CKR-1	RRPNEKK-SKAVRLIFVIMIIFFLFWTPYNLTILISVFQDFLF THECEQS
	C-C CKR-2	RCRNEKKRHRAVRVIFTIMIVYFLFWTPYNIVILLNTFQEFFGLSNCEST
	C-C CKR-3	RCPSKKK-YKAIRLIFVIMAVFFIFWTPYNVAILLSSYQSILFGND CERS
	C-C CKR-4	HCKNEKK-NKAVKMIFAVVVLFLGFWTPYNIVLFLET LVELEVLQDCTFE
15		. . . . * . . . . * . . . . . *
	M/DC CR	YNLDKSVHITKLIATTHCCINPLLYAFLDGTFSKYLRCRCH-----
	C-C CKR-1	RHLDLAVQVTEVIA YTHCCVNPVIYAFVGERFRKYLRQLFH-RRVA----
	C-C CKR-2	SQLDQATQVTETLGMTHCCINPIIYAFVGEKFRSLFHIALG-CRIAPLQK
20	C-C CKR-3	KHLDLVMLVTEVIA YSHCCMNPVIYAFVGERFRKYLRHFFH-RHLL----
	C-C CKR-4	RYLDYAIQATETLAFVHCCLNPIIYFFLGEKFRKYILQLFKTCRGLFVLC
		** . . . . * . . . . * . . . . *
	M/DC CR	-----LRSNTPLQPRGQSAQGTSREEP--DHSTEV*
25	C-C CKR-1	-----VHLVKWLPFLSVDRLERVSSTSPSTGEHEL SA---GF*
	C-C CKR-2	PVCGPGVRPGKNVKVTTQGLLDGRGKGKSIGRAPEASLQDKEGA*
	C-C CKR-3	-----MHLGRYIPFLPSEKLERTSSVSPSTAEP ELSI---VF*
	C-C CKR-4	QYCG-----LLQIYSAD-----TPSSSYTQSTMDHDLHDAL*

30

As used herein, the term "TECK" shall encompass, when used in a protein context, a protein having mature mouse or human amino acid sequences, as shown in Table 1. The invention also embraces a polypeptide comprising a

35 significant fragment of such protein. It also refers to a polypeptide which is a species counterpart, e.g., which exhibits similar biological function, and is more homologous in natural encoding sequence than other genes from that species. Typically, such chemokine will also

40 interact with its specific binding components, e.g., receptor. These binding components, e.g., antibodies, typically bind to the chemokine with high affinity, e.g., at least about 100 nM, usually better than about 30 nM, preferably better than about 10 nM, and more preferably at

45 better than about 3 nM. Homologous proteins would be found in mammalian species other than mouse, e.g., rats, dogs, cats, and primates. Non-mammalian species should also

possess structurally or functionally related genes and proteins.

The term "polypeptide" as used herein includes a significant fragment or segment, and encompasses a stretch  
5 of amino acid residues of at least about 8 amino acids, generally at least 10 amino acids, more generally at least 12 amino acids, often at least 14 amino acids, more often at least 16 amino acids, typically at least 18 amino acids, more typically at least 20 amino acids, usually at least 22  
10 amino acids, more usually at least 24 amino acids, preferably at least 26 amino acids, more preferably at least 28 amino acids, and, in particularly preferred embodiments, at least about 30 or more amino acids, e.g., about 35, 40, 45, 50, 60, 75, 80, 100, 120, etc. Similar  
15 proteins will likely comprise a plurality of such segments. Such fragments may have ends which begin and/or end at virtually all positions, e.g., beginning at residues 1, 2, 3, etc., and ending at, e.g., 69, 68, 67, 66, etc., in all combinations. Particularly interesting peptides have ends  
20 corresponding to structural domain boundaries. See, e.g., PHD and DSC programs, Rost and Sander (1994) Proteins 19:55-72; and King and Sternberg (1996) Protein Science 5:2298-2310.

The term "binding composition" refers to molecules  
25 that bind with specificity to the respective chemokine or receptor, e.g., in a ligand-receptor type fashion or an antibody-antigen interaction. These compositions may be compounds, e.g., proteins, which specifically associate with the chemokine or receptor, including natural  
30 physiologically relevant protein-protein interactions, either covalent or non-covalent. The binding composition may be a polymer, or another chemical reagent. No implication as to whether the chemokine presents a concave or convex shape in its ligand-receptor interaction is  
35 represented, other than the interaction exhibit similar specificity, e.g., specific affinity. A functional analog may be a ligand with structural modifications, or may be a

wholly unrelated molecule, e.g., which has a molecular shape which interacts with the appropriate ligand binding determinants. The ligands may serve as agonists or antagonists of the receptor, see, e.g., Goodman, et al.

- 5 (eds.) (1990) Goodman & Gilman's: The Pharmacological Bases of Therapeutics (8th ed.), Pergamon Press.

Substantially pure means that the protein is free from other contaminating proteins, nucleic acids, and/or other biologicals typically derived from the original source  
10 organism. Purity may be assayed by standard methods, and will ordinarily be at least about 40% pure, more ordinarily at least about 50% pure, generally at least about 60% pure, more generally at least about 70% pure, often at least about 75% pure, more often at least about 80% pure,  
15 typically at least about 85% pure, more typically at least about 90% pure, preferably at least about 95% pure, more preferably at least about 98% pure, and in most preferred embodiments, at least 99% pure. Analyses will typically be by weight, but may be by molar amounts.

20 Solubility of a polypeptide or fragment depends upon the environment and the polypeptide. Many parameters affect polypeptide solubility, including temperature, electrolyte environment, size and molecular characteristics of the polypeptide, and nature of the solvent. Typically,  
25 the temperature at which the polypeptide is used ranges from about 4° C to about 65° C. Usually the temperature at use is greater than about 18° C and more usually greater than about 22° C. For diagnostic purposes, the temperature will usually be about room temperature or warmer, but less  
30 than the denaturation temperature of components in the assay. For therapeutic purposes, the temperature will usually be body temperature, typically about 37° C for humans, though under certain situations the temperature may be raised or lowered in situ or in vitro.

35 The electrolytes will usually approximate in situ physiological conditions, but may be modified to higher or lower ionic strength where advantageous. The actual ions



may be modified, e.g., to conform to standard buffers used in physiological or analytical contexts.

The size and structure of the polypeptide should generally be in a substantially stable state, and usually not in a denatured state. The polypeptide may be associated with other polypeptides in a quaternary structure, e.g., to confer solubility, or associated with lipids or detergents in a manner which approximates natural lipid bilayer interactions.

The solvent will usually be a biologically compatible buffer, of a type used for preservation of biological activities, and will usually approximate a physiological solvent. Usually the solvent will have a neutral pH, typically between about 5 and 10, and preferably about 7.5. On some occasions, a detergent will be added, typically a mild non-denaturing one, e.g., CHS or CHAPS, or a low enough concentration as to avoid significant disruption of structural or physiological properties of the protein.

Solubility is reflected by sedimentation measured in Svedberg units, which are a measure of the sedimentation velocity of a molecule under particular conditions. The determination of the sedimentation velocity was classically performed in an analytical ultracentrifuge, but is typically now performed in a standard ultracentrifuge.

See, Freifelder (1982) Physical Biochemistry (2d ed.), W.H. Freeman; and Cantor and Schimmel (1980) Biophysical Chemistry, parts 1-3, W.H. Freeman & Co., San Francisco.

As a crude determination, a sample containing a putatively soluble polypeptide is spun in a standard full sized ultracentrifuge at about 50K rpm for about 10 minutes, and soluble molecules will remain in the supernatant. A soluble particle or polypeptide will typically be less than about 30S, more typically less than about 15S, usually less than about 10S, more usually less than about 6S, and, in particular embodiments, preferably less than about 4S, and more preferably less than about 3S.

### III. Physical Variants

This invention also encompasses proteins or peptides having substantial amino acid sequence homology with the amino acid sequence of each respective chemokine or receptor. The variants include species or polymorphic variants.

Amino acid sequence homology, or sequence identity, is determined by optimizing residue matches, if necessary, by introducing gaps as required. This changes when considering conservative substitutions as matches. Conservative substitutions typically include substitutions within the following groups: glycine, alanine; valine, isoleucine, leucine; aspartic acid, glutamic acid; asparagine, glutamine; serine, threonine; lysine, arginine; and phenylalanine, tyrosine. Homologous amino acid sequences are typically intended to include natural allelic and interspecies variations in each respective protein sequence. Typical homologous proteins or peptides will have from 25-100% homology (if gaps can be introduced), to 50-100% homology (if conservative substitutions are included) with the amino acid sequence of the appropriate chemokine or receptor. Homology measures will be at least about 35%, generally at least 40%, more generally at least 45%, often at least 50%, more often at least 55%, typically at least 60%, more typically at least 65%, usually at least 70%, more usually at least 75%, preferably at least 80%, and more preferably at least 80%, and in particularly preferred embodiments, at least 85% or more. See also Needleham, et al. (1970) J. Mol. Biol. 48:443-453; Sankoff, et al. (1983) Chapter One in Time Warps, String Edits, and Macromolecules: The Theory and Practice of Sequence Comparison Addison-Wesley, Reading, MA; and software packages from IntelliGenetics, Mountain View, CA; and the University of Wisconsin Genetics Computer Group, Madison, WI.

Each of the isolated chemokine or receptor DNAs can be readily modified by nucleotide substitutions, nucleotide deletions, nucleotide insertions, and inversions of nucleotide stretches. These modifications result in novel DNA sequences which encode these antigens, their derivatives, or proteins having similar physiological, immunogenic, or antigenic activity. These modified sequences can be used to produce mutant antigens or to enhance expression. Enhanced expression may involve gene amplification, increased transcription, increased translation, and other mechanisms. Such mutant chemokine or receptor derivatives include predetermined or site-specific mutations of the respective protein or its fragments. "Mutant chemokine" encompasses a polypeptide otherwise falling within the homology definition of the chemokine as set forth above, but having an amino acid sequence which differs from that of the chemokine as found in nature, whether by way of deletion, substitution, or insertion. These include substitution levels from none, one, two, three, etc. In particular, "site specific mutant chemokine" generally includes proteins having significant homology with a ligand having sequences of Table 1 through 3, and as sharing various biological activities, e.g., antigenic or immunogenic, with those sequences, and in preferred embodiments contain most of the disclosed sequences. Similar concepts apply to the different chemokine protein embodiments, particularly those found in various warm blooded animals, e.g., mammals and birds. As stated before, it is emphasized that descriptions are generally meant to encompass the various chemokine proteins, not limited to the mouse or human embodiments specifically discussed. Similar concepts apply to the receptor embodiments.

Although site specific mutation sites are often predetermined, mutants need not be site specific. Chemokine mutagenesis can be conducted by making amino acid insertions or deletions. Substitutions, deletions,

insertions, or combinations may be generated to arrive at a final construct. Insertions include amino- or carboxy-terminal fusions. Random mutagenesis can be conducted at a target codon and the expressed mutants can then be screened for the desired activity. Methods for making substitution mutations at predetermined sites in DNA having a known sequence are well known in the art, e.g., by M13 primer mutagenesis or polymerase chain reaction (PCR) techniques. See also Sambrook, et al. (1989) and Ausubel, et al. (1987 and Supplements).

The mutations in the DNA normally should not place coding sequences out of reading frames and preferably will not create complementary regions that could hybridize to produce secondary mRNA structure such as loops or hairpins.

The present invention also provides recombinant proteins, e.g., heterologous fusion proteins using segments from these proteins. A heterologous fusion protein is a fusion of proteins or segments which are naturally not normally fused in the same manner. Thus, the fusion product of an immunoglobulin with a chemokine or receptor polypeptide is a continuous protein molecule having sequences fused in a typical peptide linkage, typically made as a single translation product and exhibiting properties derived from each source peptide. A similar chimeric concept applies to heterologous nucleic acid sequences.

In addition, new constructs may be made from combining similar functional domains from other proteins. For example, ligand-binding or other segments may be "swapped" between different new fusion polypeptides or fragments. See, e.g., Cunningham, et al. (1989) Science 243:1330-1336; and O'Dowd, et al. (1988) J. Biol. Chem. 263:15985-15992. Thus, new chimeric polypeptides exhibiting new combinations of specificities will result from the functional linkage of ligand-binding specificities and other functional domains.

The phosphoramidite method described by Beaucage and Carruthers (1981) Tetra. Letts. 22:1859-1862, will produce

suitable synthetic DNA fragments. A double stranded fragment will often be obtained either by synthesizing the complementary strand and annealing the strand together under appropriate conditions or by adding the complementary strand using DNA polymerase with an appropriate primer sequence, e.g., PCR techniques.

#### IV. Functional Variants

The blocking of physiological response to various embodiments of these chemokines may result from the inhibition of binding of the ligand to its receptor, likely through competitive inhibition. Thus, in vitro assays of the present invention will often use isolated protein, membranes from cells expressing a recombinant membrane associated chemokine, soluble fragments comprising receptor binding segments of these ligands, or fragments attached to solid phase substrates. These assays will also allow for the diagnostic determination of the effects of either binding segment mutations and modifications, or ligand mutations and modifications, e.g., ligand analogs.

This invention also contemplates the use of competitive drug screening assays, e.g., where neutralizing antibodies to antigen or receptor fragments compete with a test compound for binding to the protein. In this manner, the antibodies can be used to detect the presence of polypeptides which share one or more antigenic binding sites of the ligand and can also be used to occupy binding sites on the protein that might otherwise interact with a receptor.

Additionally, neutralizing antibodies against a specific chemokine embodiment and soluble fragments of the chemokine which contain a high affinity receptor binding site, can be used to inhibit chemokine activity in tissues, e.g., tissues experiencing abnormal physiology.

"Derivatives" of chemokine antigens include amino acid sequence mutants, glycosylation variants, and covalent or aggregate conjugates with other chemical moieties.

Covalent derivatives can be prepared by linkage of functionalities to groups which are found in chemokine amino acid side chains or at the N- or C- termini, by means which are well known in the art. These derivatives can include, without limitation, aliphatic esters or amides of the carboxyl terminus, or of residues containing carboxyl side chains, O-acyl derivatives of hydroxyl group-containing residues, and N-acyl derivatives of the amino terminal amino acid or amino-group containing residues, e.g., lysine or arginine. Acyl groups are selected from the group of alkyl-moieties including C3 to C18 normal alkyl, thereby forming alkanoyl aroyl species. Covalent attachment to carrier proteins may be important when immunogenic moieties are haptens.

In particular, glycosylation alterations are included, e.g., made by modifying the glycosylation patterns of a polypeptide during its synthesis and processing, or in further processing steps. Particularly preferred means for accomplishing this are by exposing the polypeptide to glycosylating enzymes derived from cells which normally provide such processing, e.g., mammalian glycosylation enzymes. Deglycosylation enzymes are also contemplated. Also embraced are versions of the same primary amino acid sequence which have other minor modifications, including phosphorylated amino acid residues, e.g., phosphotyrosine, phosphoserine, or phosphothreonine.

A major group of derivatives are covalent conjugates of the respective chemokine or receptor or fragments thereof with other proteins or polypeptides. These derivatives can be synthesized in recombinant culture such as N- or C-terminal fusions or by the use of agents known in the art for their usefulness in cross-linking proteins through reactive side groups. Preferred chemokine derivatization sites with cross-linking agents are at free amino groups, carbohydrate moieties, and cysteine residues.

Fusion polypeptides between these chemokines and other homologous or heterologous proteins, e.g., other

chemokines, are also provided. Many growth factors and cytokines are homodimeric entities, and a repeat construct may have various advantages, including lessened susceptibility to proteolytic cleavage. Moreover, many  
5 receptors require dimerization to transduce a signal, and various dimeric ligands or domain repeats can be desirable. Homologous polypeptides may be fusions between different surface markers, resulting in, e.g., a hybrid protein exhibiting receptor binding specificity. Likewise,  
10 heterologous fusions may be constructed which would exhibit a combination of properties or activities of the derivative proteins. Typical examples are fusions of a reporter polypeptide, e.g., luciferase, with a segment or domain of a ligand, e.g., a receptor-binding segment, so that the  
15 presence or location of the fused ligand may be easily determined. See, e.g., Dull, et al., U.S. Patent No. 4,859,609. Other gene fusion partners include bacterial  $\beta$ -galactosidase, trpE, Protein A,  $\beta$ -lactamase, alpha amylase, alcohol dehydrogenase, a FLAG fusion, and yeast alpha  
20 mating factor. See, e.g., Godowski, et al. (1988) Science 241:812-816.

The phosphoramidite method described by Beaucage and Carruthers (1981) Tetra. Letts. 22:1859-1862, will produce suitable synthetic DNA fragments. A double stranded  
25 fragment will often be obtained either by synthesizing the complementary strand and annealing the strand together under appropriate conditions or by adding the complementary strand using DNA polymerase with an appropriate primer sequence.

30 Such polypeptides may also have amino acid residues which have been chemically modified by phosphorylation, sulfonation, biotinylation, or the addition or removal of other moieties, particularly those which have molecular shapes similar to phosphate groups. In some embodiments,  
35 the modifications will be useful labeling reagents, or serve as purification targets, e.g., affinity tags as FLAG.

Fusion proteins will typically be made by either recombinant nucleic acid methods or by synthetic polypeptide methods. Techniques for nucleic acid manipulation and expression are described generally, for example, in Sambrook, et al. (1989) Molecular Cloning: A Laboratory Manual (2d ed.), Vols. 1-3, Cold Spring Harbor Laboratory. Techniques for synthesis of polypeptides are described, for example, in Merrifield (1963) J. Amer. Chem. Soc. 85:2149-2156; Merrifield (1986) Science 232: 341-347; and Atherton, et al. (1989) Solid Phase Peptide Synthesis: A Practical Approach, IRL Press, Oxford; and chemical ligation, e.g., Dawson, et al. (1994) Science 266:776-779, a method of linking long synthetic peptides by a peptide bond.

This invention also contemplates the use of derivatives of these chemokines or receptors other than variations in amino acid sequence or glycosylation. Such derivatives may involve covalent or aggregative association with chemical moieties. These derivatives generally fall into the three classes: (1) salts, (2) side chain and terminal residue covalent modifications, and (3) adsorption complexes, for example with cell membranes. Such covalent or aggregative derivatives are useful as immunogens, as reagents in immunoassays, or in purification methods such as for affinity purification of ligands or other binding ligands. For example, a chemokine antigen can be immobilized by covalent bonding to a solid support such as cyanogen bromide-activated Sepharose, by methods which are well known in the art, or adsorbed onto polyolefin surfaces, with or without glutaraldehyde cross-linking, for use in the assay or purification of anti-chemokine antibodies or its receptor. These chemokines can also be labeled with a detectable group, for example radioiodinated by the chloramine T procedure, covalently bound to rare earth chelates, or conjugated to a fluorescent moiety for use in diagnostic assays. Purification of chemokine may be effected by immobilized antibodies or receptor.



Other modifications may be introduced with the goal of modifying the therapeutic pharmacokinetics or pharmacodynamics of a target chemokine. For example, certain means to minimize the size of the entity may  
5 improve its pharmacoaccessibility; other means to maximize size may affect pharmacodynamics.

A solubilized chemokine or appropriate fragment of this invention can be used as an immunogen for the production of antisera or antibodies specific for the  
10 ligand or fragments thereof. The purified chemokines can be used to screen monoclonal antibodies or chemokine-binding fragments prepared by immunization with various forms of impure preparations containing the protein. In particular, antibody equivalents include antigen binding  
15 fragments of natural antibodies, e.g., Fv, Fab, or F(ab)<sub>2</sub>. Purified chemokines can also be used as a reagent to detect antibodies generated in response to the presence of elevated levels of the protein or cell fragments containing the protein, both of which may be diagnostic of an abnormal  
20 or specific physiological or disease condition. Additionally, chemokine protein fragments, or their concatenates, may also serve as immunogens to produce antibodies of the present invention, as described immediately below. For example, this invention  
25 contemplates antibodies raised against amino acid sequences shown in Tables 1 through 3, or proteins containing them. In particular, this invention contemplates antibodies having binding affinity to or being raised against specific fragments, e.g., those which are predicted to lie on the  
30 outside surfaces of protein tertiary structure. Similar concepts apply to antibodies specific for receptors of the invention.

The present invention contemplates the isolation of additional closely related species variants. Southern and  
35 Northern blot analysis should establish that similar genetic entities exist in other mammals, and establish the stringency of hybridization conditions to isolate such. It

is likely that these chemokines and receptors are widespread in species variants, e.g., rodents, lagomorphs, carnivores, artiodactyla, perissodactyla, and primates.

The invention also provides means to isolate a group  
5 of related chemokines displaying both distinctness and similarities in structure, expression, and function. Elucidation of many of the physiological effects of the proteins will be greatly accelerated by the isolation and characterization of distinct species variants of the  
10 ligands. In particular, the present invention provides useful probes for identifying additional homologous genetic entities in different species.

The isolated genes will allow transformation of cells lacking expression of a corresponding chemokine, e.g.,  
15 either species types or cells which lack corresponding ligands and exhibit negative background activity. Expression of transformed genes will allow isolation of antigenically pure cell lines, with defined or single specie variants. This approach will allow for more  
20 sensitive detection and discrimination of the physiological effects of chemokine receptor proteins. Subcellular fragments, e.g., cytoplasts or membrane fragments, can be isolated and used.

Dissection of critical structural elements which  
25 effect the various differentiation functions provided by ligands is possible using standard techniques of modern molecular biology, particularly in comparing members of the related class. See, e.g., the homolog-scanning mutagenesis technique described in Cunningham, et al. (1989) Science  
30 243:1339-1336; and approaches used in O'Dowd, et al. (1988) J. Biol. Chem. 263:15985-15992; and Lechleiter, et al. (1990) EMBO J. 9:4381-4390.

In addition, receptor binding segments can be substituted between species variants to determine what  
35 structural features are important in both receptor binding affinity and specificity, as well as signal transduction. An array of different chemokine variants will be used to

screen for ligands exhibiting combined properties of interaction with different receptor species variants.

Intracellular functions would probably involve segments of the receptor which are normally accessible to the cytosol. However, ligand internalization may occur under certain circumstances, and interaction between intracellular components and "extracellular" segments may occur. The specific segments of interaction of a particular chemokine with other intracellular components may be identified by mutagenesis or direct biochemical means, e.g., cross-linking or affinity methods. Structural analysis by crystallographic or other physical methods will also be applicable. Further investigation of the mechanism of signal transduction will include study of associated components which may be isolatable by affinity methods or by genetic means, e.g., complementation analysis of mutants.

Further study of the expression and control of the various chemokines will be pursued. The controlling elements associated with the proteins may exhibit differential developmental, tissue specific, or other expression patterns. Upstream or downstream genetic regions, e.g., control elements, are of interest. Differential splicing of message may lead to membrane bound forms, soluble forms, and modified versions of ligand.

Structural studies of the proteins will lead to design of new ligands, particularly analogs exhibiting agonist or antagonist properties on the receptor. This can be combined with previously described screening methods to isolate ligands exhibiting desired spectra of activities.

Expression in other cell types will often result in glycosylation differences in a particular chemokine. Various species variants may exhibit distinct functions based upon structural differences other than amino acid sequence. Differential modifications may be responsible for differential function, and elucidation of the effects are now made possible.

Thus, the present invention provides important reagents related to a physiological chemokine-binding protein interaction. Although the foregoing description has focused primarily upon the mouse and human embodiments of the chemokines specifically described, those of skill in the art will immediately recognize that the invention provides other species counterparts, e.g., rat and other mammalian species or allelic or polymorphic variants.

#### 10 V. Antibodies

Antibodies can be raised to these chemokines, including species or polymorphic variants, and fragments thereof, both in their naturally occurring forms and in their recombinant forms. Additionally, antibodies can be raised to chemokines in either their active forms or in their inactive forms. Anti-idiotypic antibodies are also contemplated.

Antibodies, including binding fragments and single chain versions, against predetermined fragments of the ligands can be raised by immunization of animals with concatemers or conjugates of the fragments with immunogenic proteins. Monoclonal antibodies are prepared from cells secreting the desired antibody. These antibodies can be screened for binding to normal or defective chemokines, or screened for agonistic or antagonistic activity, e.g., mediated through a receptor for a chemokine. These monoclonal antibodies will usually bind with at least a  $K_D$  of about 1 mM, more usually at least about 300  $\mu$ M, typically at least about 10  $\mu$ M, more typically at least about 30  $\mu$ M, preferably at least about 10  $\mu$ M, and more preferably at least about 3  $\mu$ M or better.

The antibodies, including antigen binding fragments, of this invention can have significant diagnostic or therapeutic value. They can be potent antagonists that bind to ligand and inhibit binding to receptor or inhibit the ability of a ligand to elicit a biological response. They also can be useful as non-neutralizing antibodies and

can be coupled to toxins or radionuclides so that when the antibody binds to ligand, a cell expressing it, e.g., on its surface via receptor, is killed. Further, these antibodies can be conjugated to drugs or other therapeutic agents, either directly or indirectly by means of a linker, and may effect drug targeting. Antibodies to receptors may be more easily used to block ligand binding and signal transduction.

The antibodies of this invention can also be useful in diagnostic or reagent purification applications. As capture or non-neutralizing antibodies, they can be screened for ability to bind to the chemokines without inhibiting receptor binding. As neutralizing antibodies, they can be useful in competitive binding assays. They will also be useful in detecting or quantifying chemokine or, indirectly, receptors, e.g., in immunoassays. They may be used as purification reagents in immunoaffinity columns or as immunohistochemistry reagents.

Ligand fragments may be concatenated or joined to other materials, particularly polypeptides, as fused or covalently joined polypeptides to be used as immunogens. Short peptides will preferably be made as repeat structures to increase size. A ligand and its fragments may be fused or covalently linked to a variety of immunogens, such as keyhole limpet hemocyanin, bovine serum albumin, tetanus toxoid, etc. See Microbiology, Hoeber Medical Division, Harper and Row, 1969; Landsteiner (1962) Specificity of Serological Reactions, Dover Publications, New York, and Williams, et al. (1967) Methods in Immunology and Immunochemistry, Vol. 1, Academic Press, New York, for descriptions of methods of preparing polyclonal antisera. A typical method involves hyperimmunization of an animal with an antigen. The blood of the animal is then collected shortly after the repeated immunizations and the gamma globulin fraction is isolated.

In some instances, it is desirable to prepare monoclonal antibodies from various mammalian hosts, such as

mice, rodents, primates, humans, etc. Description of techniques for preparing such monoclonal antibodies may be found in, e.g., Stites, et al. (eds.) Basic and Clinical Immunology (4th ed.), Lange Medical Publications, Los Altos, CA, and references cited therein; Harlow and Lane (1988) Antibodies: A Laboratory Manual, CSH Press; Goding (1986) Monoclonal Antibodies: Principles and Practice (2d ed.) Academic Press, New York; and particularly in Kohler and Milstein (1975) in Nature 256:495-497, which discusses one method of generating monoclonal antibodies. Summarized briefly, this method involves injecting an animal with an immunogen. The animal is then sacrificed and cells taken, e.g., from its spleen, which are then fused with myeloma cells. The result is a hybrid cell or "hybridoma" that is capable of reproducing in vitro. The population of hybridomas is then screened to isolate individual clones, each of which secrete a single antibody species to the immunogen. In this manner, the individual antibody species obtained are the products of immortalized and cloned single B cells from the immune animal generated in response to a specific site recognized on the immunogenic substance. Large amounts of antibody may be derived from ascites fluid from an animal.

Other suitable techniques involve in vitro exposure of lymphocytes to the antigenic polypeptides or alternatively to selection of libraries of antibodies in phage or similar vectors. See, Huse, et al. (1989) "Generation of a Large Combinatorial Library of the Immunoglobulin Repertoire in Phage Lambda," Science 246:1275-1281; and Ward, et al. (1989) Nature 341:544-546. The polypeptides and antibodies of the present invention may be used with or without modification, including chimeric or humanized antibodies. Frequently, the polypeptides and antibodies will be labeled by joining, either covalently or non-covalently, a substance which provides for a detectable signal. A wide variety of labels and conjugation techniques are known and are reported extensively in both the scientific and patent

literature. Suitable labels include radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent moieties, chemiluminescent moieties, magnetic particles, and the like. Patents, teaching the use of such labels include U.S. Patent Nos. 3,817,837; 3,850,752; 3,939,350; 3,996,345; 4,277,437; 4,275,149; and 4,366,241. Also, recombinant immunoglobulins may be produced, see Cabilly, U.S. Patent No. 4,816,567; and Queen et al. (1989) Proc. Nat'l. Acad. Sci. 86:10029-10033.

10       The antibodies of this invention can also be used for affinity chromatography in isolating the protein. Columns can be prepared where the antibodies are linked to a solid support, e.g., particles, such as agarose, Sephadex, or the like, where a cell lysate may be passed through the column, 15 the column washed, followed by increasing concentrations of a mild denaturant, whereby the purified chemokine protein will be released.

      The antibodies may also be used to screen expression libraries for particular expression products. Usually the 20 antibodies used in such a procedure will be labeled with a moiety allowing easy detection of presence of antigen by antibody binding.

      Antibodies raised against these chemokine will also be useful to raise anti-idiotypic antibodies. These will be 25 useful in detecting or diagnosing various immunological conditions related to expression of the respective antigens.

#### VI. Nucleic Acids

30       The described peptide sequences and the related reagents are useful in isolating a DNA clone encoding these chemokines, e.g., from a natural source. Typically, it will be useful in isolating a gene from another individual, and similar procedures will be applied to isolate genes 35 from other species, e.g., warm blooded animals, such as birds and mammals. Cross hybridization will allow isolation of ligand from other species. A number of

different approaches should be available to successfully isolate a suitable nucleic acid clone. Similar concepts apply to the receptor embodiments.

The purified protein or defined peptides are useful  
5 for generating antibodies by standard methods, as described above. Synthetic peptides or purified protein can be presented to an immune system to generate monoclonal or polyclonal antibodies. See, e.g., Coligan (1991) Current Protocols in Immunology Wiley/Greene; and Harlow and Lane  
10 (1989) Antibodies: A Laboratory Manual Cold Spring Harbor Press. Alternatively, a chemokine receptor can be used as a specific binding reagent, and advantage can be taken of its specificity of binding, much like an antibody would be used. However, chemokine receptors are typically 7  
15 transmembrane proteins, which could be sensitive to appropriate interaction with lipid or membrane. The signal transduction typically is mediated through a G-protein.

For example, the specific binding composition could be used for screening of an expression library made from a  
20 cell line which expresses a particular chemokine. The screening can be standard staining of surface expressed ligand, or by panning. Screening of intracellular expression can also be performed by various staining or immunofluorescence procedures. The binding compositions  
25 could be used to affinity purify or sort out cells expressing the ligand.

The peptide segments can also be used to predict appropriate oligonucleotides to screen a library, e.g., to isolate species variants. The genetic code can be used to  
30 select appropriate oligonucleotides useful as probes for screening. See, e.g., Tables 1 through 5. In combination with polymerase chain reaction (PCR) techniques, synthetic oligonucleotides will be useful in selecting correct clones from a library. Complementary sequences will also be used  
35 as probes or primers. Based upon identification of the likely amino terminus, the third peptide should be particularly useful, e.g., coupled with anchored vector or



poly-A complementary PCR techniques or with complementary DNA of other peptides.

This invention contemplates use of isolated DNA or fragments to encode a biologically active corresponding chemokine polypeptide. In addition, this invention covers isolated or recombinant DNA which encodes a biologically active protein or polypeptide which is capable of hybridizing under appropriate conditions with the DNA sequences described herein. Said biologically active protein or polypeptide can be an intact ligand, or fragment, and have an amino acid sequence as disclosed in Tables 1 through 3. Further, this invention covers the use of isolated or recombinant DNA, or fragments thereof, which encode proteins which are homologous to a chemokine or which was isolated using cDNA encoding a chemokine as a probe. The isolated DNA can have the respective regulatory sequences in the 5' and 3' flanks, e.g., promoters, enhancers, poly-A addition signals, and others. Alternatively, promoters or other regulatory signals may be incorporated to be operably linked to natural genes in a cell.

An "isolated" nucleic acid is a nucleic acid, e.g., an RNA, DNA, or a mixed polymer, which is substantially separated from other components which naturally accompany a native sequence, e.g., ribosomes, polymerases, and flanking genomic sequences from the originating species. The term embraces a nucleic acid sequence which has been removed from its naturally occurring environment, and includes recombinant or cloned DNA isolates and chemically synthesized analogs or analogs biologically synthesized by heterologous systems. A substantially pure molecule includes isolated forms of the molecule.

An isolated nucleic acid will generally be a homogeneous composition of molecules, but will, in some embodiments, contain minor heterogeneity. This heterogeneity is typically found at the polymer ends or

portions not critical to a desired biological function or activity.

A "recombinant" nucleic acid is defined either by its method of production or its structure. In reference to its method of production, e.g., a product made by a process, the process is use of recombinant nucleic acid techniques, e.g., involving human intervention in the nucleotide sequence, typically selection or production. Alternatively, it can be a nucleic acid made by generating a sequence comprising fusion of two fragments which are not naturally contiguous to each other, but is meant to exclude products of nature, e.g., naturally occurring mutants. Thus, for example, products made by transforming cells with any unnaturally occurring vector is encompassed, as are nucleic acids comprising sequence derived using any synthetic oligonucleotide process. Such is often done to replace a codon with a redundant codon encoding the same or a conservative amino acid, while typically introducing or removing a sequence recognition site. Alternatively, it is performed to join together nucleic acid segments of desired functions to generate a single genetic entity comprising a desired combination of functions not found in the commonly available natural forms. Restriction enzyme recognition sites are often the target of such artificial manipulations, but other site specific targets, e.g., promoters, DNA replication sites, regulation sequences, control sequences, or other useful features may be incorporated by design. A similar concept is intended for a recombinant, e.g., fusion, polypeptide. Specifically included are synthetic nucleic acids which, by genetic code redundancy, encode polypeptides similar to fragments of these antigens, and fusions of sequences from various different species variants.

A significant "fragment" in a nucleic acid context is a contiguous segment of at least about 17 nucleotides, generally at least about 20 nucleotides, more generally at least about 23 nucleotides, ordinarily at least about 26

nucleotides, more ordinarily at least about 29 nucleotides, often at least about 32 nucleotides, more often at least about 35 nucleotides, typically at least about 38 nucleotides, more typically at least about 41 nucleotides, usually at least about 44 nucleotides, more usually at least about 47 nucleotides, preferably at least about 50 nucleotides, more preferably at least about 53 nucleotides, and in particularly preferred embodiments will be at least about 56 or more nucleotides, e.g., 60, 65, 75, 85, 100, 120, 150, 200, 250, 300, 400, etc. Such fragments may have ends which begin and/or end at virtually all positions, e.g., beginning at nucleotides 1, 2, 3, etc., and ending at, e.g., 300, 299, 298, 287, etc., in all combinations. Particularly interesting polynucleotides have ends corresponding to structural domain boundaries.

A DNA which codes for a particular chemokine protein or peptide will be very useful to identify genes, mRNA, and cDNA species which code for related or homologous ligands, as well as DNAs which code for homologous proteins from different species. There are likely homologs in other species, including primates. Various chemokine proteins should be homologous and are encompassed herein. However, even proteins that have a more distant evolutionary relationship to the ligand can readily be isolated under appropriate conditions using these sequences if they are sufficiently homologous. Primate chemokines are of particular interest.

This invention further covers recombinant DNA molecules and fragments having a DNA sequence identical to or highly homologous to the isolated DNAs set forth herein. In particular, the sequences will often be operably linked to DNA segments which control transcription, translation, and DNA replication. Alternatively, recombinant clones derived from the genomic sequences, e.g., containing introns, will be useful for transgenic studies, including, e.g., transgenic cells and organisms, and for gene therapy. See, e.g., Goodnow (1992) "Transgenic Animals" in Roitt

- (ed.) Encyclopedia of Immunology Academic Press, San Diego, pp. 1502-1504; Travis (1992) Science 256:1392-1394; Kuhn, et al. (1991) Science 254:707-710; Capecchi (1989) Science 244:1288; Robertson (1987) (ed.) Teratocarcinomas and  
5 Embryonic Stem Cells: A Practical Approach IRL Press, Oxford; and Rosenberg (1992) J. Clinical Oncology 10:180-199.

Homologous nucleic acid sequences, when compared, exhibit significant similarity, or identity. The standards  
10 for homology in nucleic acids are either measures for homology generally used in the art by sequence comparison or based upon hybridization conditions. The hybridization conditions are described in greater detail below.

Substantial homology in the nucleic acid sequence  
15 comparison context means either that the segments, or their complementary strands, when compared, are identical when optimally aligned, with appropriate nucleotide insertions or deletions, in at least about 50% of the nucleotides, generally at least about 56%, more generally at least about  
20 59%, ordinarily at least about 62%, more ordinarily at least about 65%, often at least about 68%, more often at least about 71%, typically at least about 74%, more typically at least about 77%, usually at least about 80%, more usually at least about 85%, preferably at least about  
25 90%, more preferably at least about 95 to 98% or more, and in particular embodiments, as high at about 99% or more of the nucleotides. Alternatively, substantial homology exists when the segments will hybridize under selective hybridization conditions, to a strand, or its complement,  
30 typically using a sequence derived from Tables 1 through 5. Typically, selective hybridization will occur when there is at least about 55% homology over a stretch of at least about 30 nucleotides, preferably at least about 65% over a stretch of at least about 25 nucleotides, more preferably  
35 at least about 75%, and most preferably at least about 90% over about 20 nucleotides. See, Kanehisa (1984) Nuc. Acids Res. 12:203-213. The length of homology comparison, as

described, may be over longer stretches, and in certain  
embodiments will be over a stretch of at least about 17  
nucleotides, usually at least about 20 nucleotides, more  
usually at least about 24 nucleotides, typically at least  
5 about 28 nucleotides, more typically at least about 40  
nucleotides, preferably at least about 50 nucleotides, and  
more preferably at least about 75 to 100 or more  
nucleotides.

Stringent conditions, in referring to homology in the  
10 hybridization context, will be stringent combined  
conditions of salt, temperature, organic solvents, and  
other parameters, typically those controlled in  
hybridization reactions. Stringent temperature conditions  
will usually include temperatures in excess of about 30° C,  
15 more usually in excess of about 37° C, typically in excess  
of about 45° C, more typically in excess of about 55° C,  
preferably in excess of about 65° C, and more preferably in  
excess of about 70° C. Stringent salt conditions will  
ordinarily be less than about 1000 mM, usually less than  
20 about 500 mM, more usually less than about 400 mM,  
typically less than about 300 mM, preferably less than  
about 200 mM, and more preferably less than about 150 mM.  
However, the combination of parameters is much more  
important than the measure of any single parameter. See,  
25 e.g., Wetmur and Davidson (1968) J. Mol. Biol. 31:349-370.

Corresponding chemokines from other mammalian species  
can be cloned and isolated by cross-species hybridization  
of closely related species. Alternatively, sequences from  
a data base may be recognized as having similarity.  
30 Homology may be relatively low between distantly related  
species, and thus hybridization of relatively closely  
related species is advisable. Alternatively, preparation  
of an antibody preparation which exhibits less species  
specificity may be useful in expression cloning approaches.  
35 PCR approaches using segments of conserved sequences will  
also be used.

## VII. Making chemokines, receptors; Mimetics

DNA which encodes each respective chemokine or fragments thereof can be obtained by chemical synthesis, screening cDNA libraries, or by screening genomic libraries prepared from a wide variety of cell lines or tissue samples.

This DNA can be expressed in a wide variety of host cells for the synthesis of a full-length ligand or fragments which can in turn, for example, be used to generate polyclonal or monoclonal antibodies; for binding studies; for construction and expression of modified molecules; and for structure/function studies. Each antigen or its fragments can be expressed in host cells that are transformed or transfected with appropriate expression vectors. These molecules can be substantially purified to be free of protein or cellular contaminants, other than those derived from the recombinant host, and therefore are particularly useful in pharmaceutical compositions when combined with a pharmaceutically acceptable carrier and/or diluent. The antigen, or portions thereof, may be expressed as fusions with other proteins.

Expression vectors are typically self-replicating DNA or RNA constructs containing the desired antigen gene or its fragments, usually operably linked to suitable genetic control elements that are recognized in a suitable host cell. These control elements are capable of effecting expression within a suitable host. The specific type of control elements necessary to effect expression will depend upon the eventual host cell used. Generally, the genetic control elements can include a prokaryotic promoter system or a eukaryotic promoter expression control system, and typically include a transcriptional promoter, an optional operator to control the onset of transcription, transcription enhancers to elevate the level of mRNA expression, a sequence that encodes a suitable ribosome binding site, and sequences that terminate transcription.

and translation. Expression vectors also usually contain an origin of replication that allows the vector to replicate independently of the host cell.

The vectors of this invention contain DNA which encode  
5   embodiments of a chemokine, receptor, or a fragment  
     thereof, typically encoding a biologically active  
     polypeptide. The DNA can be under the control of a viral  
     promoter and can encode a selection marker. This invention  
     further contemplates use of such expression vectors which  
10   are capable of expressing eukaryotic cDNA coding for each  
     chemokine in a prokaryotic or eukaryotic host, where the  
     vector is compatible with the host and where the eukaryotic  
     cDNA coding for the ligand is inserted into the vector such  
     that growth of the host containing the vector expresses the  
15   cDNA in question. Usually, expression vectors are designed  
     for stable replication in their host cells or for  
     amplification to greatly increase the total number of  
     copies of the desirable gene per cell. It is not always  
     necessary to require that an expression vector replicate in  
20   a host cell, e.g., it is possible to effect transient  
     expression of the ligand or its fragments in various hosts  
     using vectors that do not contain a replication origin that  
     is recognized by the host cell. It is also possible to use  
     vectors that cause integration of a chemokine gene or its  
25   fragments into the host DNA by recombination, or to  
     integrate a promoter which controls expression of an  
     endogenous gene.

Vectors, as used herein, comprise plasmids, viruses,  
bacteriophage, integratable DNA fragments, and other  
30   vehicles which enable the integration of DNA fragments into  
     the genome of the host. Expression vectors are specialized  
     vectors which contain genetic control elements that effect  
     expression of operably linked genes. Plasmids are the most  
     commonly used form of vector but all other forms of vectors  
35   which serve an equivalent function and which are, or  
     become, known in the art are suitable for use herein. See,  
     e.g., Pouwels, et al. (1985 and Supplements) Cloning

Vectors: A Laboratory Manual, Elsevier, N.Y., and  
Rodriguez, et al. (1988) (eds.) Vectors: A Survey of  
Molecular Cloning Vectors and Their Uses, Butterworth,  
Boston, MA.

5 Transformed cells include cells, preferably mammalian,  
that have been transformed or transfected with a chemokine  
gene containing vector constructed using recombinant DNA  
techniques. Transformed host cells usually express the  
ligand or its fragments, but for purposes of cloning,  
10 amplifying, and manipulating its DNA, do not need to  
express the protein. This invention further contemplates  
culturing transformed cells in a nutrient medium, thus  
permitting the protein to accumulate in the culture. The  
protein can be recovered, either from the culture or from  
15 the culture medium.

For purposes of this invention, DNA sequences are  
operably linked when they are functionally related to each  
other. For example, DNA for a presequence or secretory  
leader is operably linked to a polypeptide if it is  
20 expressed as a preprotein or participates in directing the  
polypeptide to the cell membrane or in secretion of the  
polypeptide. A promoter is operably linked to a coding  
sequence if it controls the transcription of the  
polypeptide; a ribosome binding site is operably linked to  
25 a coding sequence if it is positioned to permit  
translation. Usually, operably linked means contiguous and  
in reading frame, however, certain genetic elements such as  
repressor genes are not contiguously linked but still bind  
to operator sequences that in turn control expression.

30 Suitable host cells include prokaryotes, lower  
eukaryotes, and higher eukaryotes. Prokaryotes include  
both gram negative and gram positive organisms, e.g., E.  
coli and B. subtilis. Lower eukaryotes include yeasts,  
e.g., S. cerevisiae and Pichia, and species of the genus  
35 Dictyostelium. Higher eukaryotes include established  
tissue culture cell lines from animal cells, both of



non-mammalian origin, e.g., insect cells, and birds, and of mammalian origin, e.g., human, primates, and rodents.

Prokaryotic host-vector systems include a wide variety of vectors for many different species. As used herein, E. coli and its vectors will be used generically to include equivalent vectors used in other prokaryotes. A representative vector for amplifying DNA is pBR322 or many of its derivatives. Vectors that can be used to express these chemokines or their fragments include, but are not limited to, such vectors as those containing the lac promoter (pUC-series); trp promoter (pBR322-trp); Ipp promoter (the pIN-series); lambda-pP or pR promoters (pOTS); or hybrid promoters such as ptac (pDR540). See Brosius, et al. (1988) "Expression Vectors Employing Lambda-, trp-, lac-, and Ipp-derived Promoters", in Rodriguez and Denhardt (eds.) Vectors: A Survey of Molecular Cloning Vectors and Their Uses, Butterworth, Boston, Chapter 10, pp. 205-236.

Lower eukaryotes, e.g., yeasts and Dictyostelium, may be transformed with chemokine sequence containing vectors. For purposes of this invention, the most common lower eukaryotic host is the baker's yeast, *Saccharomyces cerevisiae*. It will be used to generically represent lower eukaryotes although a number of other strains and species are also available. Yeast vectors typically consist of a replication origin (unless of the integrating type), a selection gene, a promoter, DNA encoding the desired protein or its fragments, and sequences for translation termination, polyadenylation, and transcription termination. Suitable expression vectors for yeast include such constitutive promoters as 3-phosphoglycerate kinase and various other glycolytic enzyme gene promoters or such inducible promoters as the alcohol dehydrogenase 2 promoter or metallothionine promoter. Suitable vectors include derivatives of the following types: self-replicating low copy number (such as the YRp-series), self-replicating high copy number (such as the YEpl-series); integrating types

(such as the YIp-series), or mini-chromosomes (such as the YCp-series).

Higher eukaryotic tissue culture cells are the preferred host cells for expression of the functionally active chemokine protein. In principle, most any higher eukaryotic tissue culture cell line is workable, e.g., insect baculovirus expression systems, whether from an invertebrate or vertebrate source. However, mammalian cells are preferred, in that the processing, both cotranslationally and posttranslationally. Transformation or transfection and propagation of such cells has become a routine procedure. Examples of useful cell lines include HeLa cells, Chinese hamster ovary (CHO) cell lines, baby rat kidney (BRK) cell lines, insect cell lines, bird cell lines, and monkey (COS) cell lines. Expression vectors for such cell lines usually include an origin of replication, a promoter, a translation initiation site, RNA splice sites (if genomic DNA is used), a polyadenylation site, and a transcription termination site. These vectors also usually contain a selection gene or amplification gene. Suitable expression vectors may be plasmids, viruses, or retroviruses carrying promoters derived, e.g., from such sources as from adenovirus, SV40, parvoviruses, vaccinia virus, or cytomegalovirus. Representative examples of suitable expression vectors include pCDNA1; pCD, see Okayama, et al. (1985) Mol. Cell Biol. 5:1136-1142; pMC1neo Poly-A, see Thomas, et al. (1987) Cell 51:503-512; and a baculovirus vector such as pAC 373 or pAC 610.

It will often be desired to express a chemokine polypeptide in a system which provides a specific or defined glycosylation pattern. In this case, the usual pattern will be that provided naturally by the expression system. However, the pattern will be modifiable by exposing the polypeptide, e.g., an unglycosylated form, to appropriate glycosylating proteins introduced into a heterologous expression system. For example, a chemokine gene may be co-transformed with one or more genes encoding

mammalian or other glycosylating enzymes. Using this approach, certain mammalian glycosylation patterns will be achievable or approximated in prokaryote or other cells.

5 A chemokine, or a fragment thereof, may be engineered to be phosphatidyl inositol (PI) linked to a cell membrane, but can be removed from membranes by treatment with a phosphatidyl inositol cleaving enzyme, e.g., phosphatidyl inositol phospholipase-C. This releases the antigen in a biologically active form, and allows purification by  
10 standard procedures of protein chemistry. See, e.g., Low (1989) Biochim. Biophys. Acta 988:427-454; Tse, et al. (1985) Science 230:1003-1008; and Brunner, et al. (1991) J. Cell Biol. 114:1275-1283.

Now that these chemokines have been characterized,  
15 fragments or derivatives thereof can be prepared by conventional processes for synthesizing peptides. These include processes such as are described in Stewart and Young (1984) Solid Phase Peptide Synthesis, Pierce Chemical Co., Rockford, IL; Bodanszky and Bodanszky (1984) The Practice of Peptide Synthesis, Springer-Verlag, New York;  
20 and Bodanszky (1984) The Principles of Peptide Synthesis, Springer-Verlag, New York. For example, an azide process, an acid chloride process, an acid anhydride process, a mixed anhydride process, an active ester process (for  
25 example, p-nitrophenyl ester, N-hydroxysuccinimide ester, or cyanomethyl ester), a carbodiimidazole process, an oxidative-reductive process, or a dicyclohexylcarbodiimide (DCCD)/additive process can be used. Solid phase and solution phase syntheses are both applicable to the  
30 foregoing processes.

These chemokines, fragments, or derivatives are suitably prepared in accordance with the above processes as typically employed in peptide synthesis, generally either  
35 by a so-called stepwise process which comprises condensing an amino acid to the terminal amino acid, one by one in sequence, or by coupling peptide fragments to the terminal amino acid. Amino groups that are not being used in the

coupling reaction are typically protected to prevent coupling at an incorrect location.

If a solid phase synthesis is adopted, the C-terminal amino acid is typically bound to an insoluble carrier or support through its carboxyl group. The insoluble carrier is not particularly limited as long as it has a binding capability to a reactive carboxyl group. Examples of such insoluble carriers include halomethyl resins, such as chloromethyl resin or bromomethyl resin, hydroxymethyl resins, phenol resins, tert-alkyloxycarbonyl-hydrazidated resins, and the like.

An amino group-protected amino acid is bound in sequence through condensation of its activated carboxyl group and the reactive amino group of the previously formed peptide or chain, to synthesize the peptide step by step. After synthesizing the complete sequence, the peptide is split off from the insoluble carrier to produce the peptide. This solid-phase approach is generally described by Merrifield, et al. (1963) in J. Am. Chem. Soc. 85:2149-2156.

The prepared ligand and fragments thereof can be isolated and purified from the reaction mixture by means of peptide separation, e.g., by extraction, precipitation, electrophoresis and various forms of chromatography, and the like. The various chemokines of this invention can be obtained in varying degrees of purity depending upon its desired use. Purification can be accomplished by use of the protein purification techniques disclosed herein or by the use of the antibodies herein described, e.g., in immunoabsorbant affinity chromatography. This immunoabsorbant affinity chromatography is carried out by first linking the antibodies to a solid support and then contacting the linked antibodies with solubilized lysates of appropriate source cells, lysates of other cells expressing the ligand, or lysates or supernatants of cells producing the desired chemokine as a result of DNA techniques, see below.

## VIII. Uses

The present invention provides reagents which will find use in diagnostic applications as described elsewhere herein, e.g., in the general description for developmental abnormalities, or below in the description of kits for diagnosis.

This invention also provides reagents with significant therapeutic value. These chemokines (naturally occurring or recombinant), fragments thereof and antibodies thereto, along with compounds identified as having binding affinity to them, should be useful in the treatment of conditions associated with abnormal physiology or development, including inflammatory conditions, including asthma. In particular, modulation of trafficking of leukocytes is one likely biological activity, but a wider tissue distribution might suggest broader biological activity, including, e.g., antiviral effects. Abnormal proliferation, regeneration, degeneration, and atrophy may be modulated by appropriate therapeutic treatment using the compositions provided herein. For example, a disease or disorder associated with abnormal expression or abnormal signaling by a chemokine should be a likely target for an agonist or antagonist of the ligand.

Various abnormal physiological or developmental conditions are known in cell types shown to possess the chemokine mRNAs by Northern blot analysis. See Berkow (ed.) The Merck Manual of Diagnosis and Therapy, Merck & Co., Rahway, N.J.; and Thorn, et al. Harrison's Principles of Internal Medicine, McGraw-Hill, N.Y. Developmental or functional abnormalities, e.g., of the immune system, cause significant medical abnormalities and conditions which may be susceptible to prevention or treatment using compositions provided herein.

Chemokine antibodies, including recombinant forms, can be purified and then administered to a patient. These reagents can be combined for therapeutic use with

additional active or inert ingredients, e.g., in conventional pharmaceutically acceptable carriers or diluents, e.g., immunogenic adjuvants, along with physiologically innocuous stabilizers and excipients.

- 5 These combinations can be sterile filtered and placed into dosage forms as by lyophilization in dosage vials or storage in stabilized aqueous preparations. This invention also contemplates use of antibodies or binding fragments thereof, including forms which are not complement binding.
- 10 Moreover, modifications to the antibody molecules or antigen binding fragments thereof, may be adopted which affect the pharmacokinetics or pharmacodynamics of the therapeutic entity.

- Drug screening using antibodies or receptor or
- 15 fragments thereof can be performed to identify compounds having binding affinity to each chemokine or receptor, including isolation of associated components. Subsequent biological assays can then be utilized to determine if the compound has intrinsic stimulating activity and is
- 20 therefore a blocker or antagonist in that it blocks the activity of the ligand. Likewise, a compound having intrinsic stimulating activity can activate the receptor and is thus an agonist in that it simulates the activity of a chemokine. This invention further contemplates the
- 25 therapeutic use of antibodies to these chemokines as antagonists. This approach should be particularly useful with other chemokine species variants.

- The quantities of reagents necessary for effective therapy will depend upon many different factors, including
- 30 means of administration, target site, physiological state of the patient, and other medicants administered. Thus, treatment dosages should be titrated to optimize safety and efficacy in various populations, including racial subgroups, age, gender, etc. Typically, dosages used in
- 35 vitro may provide useful guidance in the amounts useful for in situ administration of these reagents. Animal testing of effective doses for treatment of particular disorders

will provide further predictive indication of human dosage. Various considerations are described, e.g., in Gilman, et al. (eds.) (1990) Goodman and Gilman's: The Pharmacological Bases of Therapeutics, 8th Ed., Pergamon Press; and Remington's Pharmaceutical Sciences, 17th ed. (1990), Mack Publishing Co., Easton, Penn.. Methods for administration are discussed therein and below, e.g., for oral, intravenous, intraperitoneal, or intramuscular administration, transdermal diffusion, and others.

Pharmaceutically acceptable carriers typically include water, saline, buffers, and other compounds described, e.g., in the Merck Index, Merck & Co., Rahway, New Jersey. Dosage ranges would ordinarily be expected to be in amounts lower than 1 mM concentrations, typically less than about 10  $\mu$ M concentrations, usually less than about 100 nM, preferably less than about 10 pM (picomolar), and most preferably less than about 1 fM (femtomolar), with an appropriate carrier. Slow release formulations, or a slow release apparatus will often be utilized for continuous administration.

A chemokine, fragments thereof, or antibodies to it or its fragments, antagonists, and agonists, may be administered directly to the host to be treated or, depending on the size of the compounds, it may be desirable to conjugate them to carrier proteins such as ovalbumin or serum albumin prior to their administration. Therapeutic formulations may be administered in any conventional dosage formulation. While it is possible for the active ingredient to be administered alone, it is often preferable to present it as a pharmaceutical formulation.

Formulations typically comprise at least one active ingredient, as defined above, together with one or more acceptable carriers thereof. Each carrier should be both pharmaceutically and physiologically acceptable in the sense of being compatible with the other ingredients and not injurious to the patient. Carriers may improve storage life, stability, etc. Formulations include those suitable

for oral, rectal, nasal, or parenteral (including subcutaneous, intramuscular, intravenous and intradermal) administration. The formulations may conveniently be presented in unit dosage form and may be prepared by any methods well known in the art of pharmacy. See, e.g., Gilman, et al. (eds.) (1990) Goodman and Gilman's: The Pharmacological Bases of Therapeutics, 8th Ed., Pergamon Press; and Remington's Pharmaceutical Sciences, 17th ed. (1990), Mack Publishing Co., Easton, Penn.; Avis, et al. (eds.) (1993) Pharmaceutical Dosage Forms: Parenteral Medications Dekker, New York; Lieberman, et al. (eds.) (1990) Pharmaceutical Dosage Forms: Tablets Dekker, New York; and Lieberman, et al. (eds.) (1990) Pharmaceutical Dosage Forms: Disperse Systems Dekker, New York. The therapy of this invention may be combined with or used in association with other therapeutic agents.

Both the naturally occurring and the recombinant forms of the chemokines of this invention are particularly useful in kits and assay methods which are capable of screening compounds for binding activity to the proteins. Several methods of automating assays have been developed in recent years so as to permit screening of tens of thousands of compounds in a short period. See, e.g., Fodor, et al. (1991) Science 251:767-773, which describes means for testing of binding affinity by a plurality of defined polymers synthesized on a solid substrate. The development of suitable assays can be greatly facilitated by the availability of large amounts of purified, soluble chemokine as provided by this invention.

For example, antagonists can normally be found once the ligand has been structurally defined. Testing of potential ligand analogs is now possible upon the development of highly automated assay methods using physiologically responsive cells. In particular, new agonists and antagonists will be discovered by using screening techniques described herein.



Viabile cells could also be used to screen for the effects of drugs on respective chemokine mediated functions, e.g., second messenger levels, i.e.,  $\text{Ca}^{++}$ ; inositol phosphate pool changes (see, e.g., Berridge (1993) Nature 361:315-325 or Billah and Anthes (1990) Biochem. J. 269:281-291); cellular morphology modification responses; phosphoinositide lipid turnover; an antiviral response. and others. Some detection methods allow for elimination of a separation step, e.g., a proximity sensitive detection system. Calcium sensitive dyes will be useful for detecting  $\text{Ca}^{++}$  levels, with a fluorimeter or a fluorescence cell sorting apparatus.

Rational drug design may also be based upon structural studies of the molecular shapes of the chemokines and other effectors or analogs. Effectors may be other proteins which mediate other functions in response to ligand binding, or other proteins which normally interact with the receptor. One means for determining which sites interact with specific other proteins is a physical structure determination, e.g., x-ray crystallography or 2 dimensional NMR techniques. These will provide guidance as to which amino acid residues form molecular contact regions. For a detailed description of protein structural determination, see, e.g., Blundell and Johnson (1976) Protein Crystallography, Academic Press, New York.

Purified chemokine can be coated directly onto plates for use in the aforementioned drug screening techniques. However, non-neutralizing antibodies to these ligands can be used as capture antibodies to immobilize the respective ligand on the solid phase.

Similar concepts also apply to the chemokine receptor embodiments of the invention.

#### IX. Kits

This invention also contemplates use of chemokine proteins, fragments thereof, peptides, and their fusion products in a variety of diagnostic kits and methods for

detecting the presence of ligand, antibodies, or chemokine receptors. Typically the kit will have a compartment containing either a defined chemokine peptide or gene segment or a reagent which recognizes one or the other,  
5 e.g., antibodies.

A kit for determining the binding affinity of a test compound to a chemokine would typically comprise a test compound; a labeled compound, for example an antibody having known binding affinity for the ligand; a source of  
10 chemokine (naturally occurring or recombinant); and a means for separating bound from free labeled compound, such as a solid phase for immobilizing the ligand. Once compounds are screened, those having suitable binding affinity to the ligand can be evaluated in suitable biological assays, as  
15 are well known in the art, to determine whether they act as agonists or antagonists to the receptor. The availability of recombinant chemokine polypeptides also provide well defined standards for calibrating such assays or as positive control samples.

A preferred kit for determining the concentration of, for example, a chemokine in a sample would typically comprise a labeled compound, e.g., antibody, having known binding affinity for the ligand, a source of ligand (naturally occurring or recombinant) and a means for  
20 separating the bound from free labeled compound, for example, a solid phase for immobilizing the chemokine. Compartments containing reagents, and instructions for use or disposal, will normally be provided.

Antibodies, including antigen binding fragments,  
30 specific for the chemokine or ligand fragments are useful in diagnostic applications to detect the presence of elevated levels of chemokine and/or its fragments. Such diagnostic assays can employ lysates, live cells, fixed cells, immunofluorescence, cell cultures, body fluids, and  
35 further can involve the detection of antigens related to the ligand in serum, or the like. Diagnostic assays may be homogeneous (without a separation step between free reagent

and antigen-ligand complex) or heterogeneous (with a separation step). Various commercial assays exist, such as radioimmunoassay (RIA), enzyme-linked immunosorbent assay (ELISA), enzyme immunoassay (EIA), enzyme-multiplied  
5 immunoassay technique (EMIT), substrate-labeled fluorescent immunoassay (SLFIA), and the like. For example, unlabeled antibodies can be employed by using a second antibody which is labeled and which recognizes the antibody to a chemokine or to a particular fragment thereof. Similar assays have  
10 also been extensively discussed in the literature. See, e.g., Harlow and Lane (1988) Antibodies: A Laboratory Manual, CSH.

Anti-idiotypic antibodies may have similar uses to diagnose presence of antibodies against a chemokine, as  
15 such may be diagnostic of various abnormal states. For example, overproduction of a chemokine may result in production of various immunological reactions which may be diagnostic of abnormal physiological states, particularly in various inflammatory or asthma conditions.

20 Frequently, the reagents for diagnostic assays are supplied in kits, so as to optimize the sensitivity of the assay. For the subject invention, depending upon the nature of the assay, the protocol, and the label, either labeled or unlabeled antibody or labeled chemokine is  
25 provided. This is usually in conjunction with other additives, such as buffers, stabilizers, materials necessary for signal production such as substrates for enzymes, and the like. Preferably, the kit will also contain instructions for proper use and disposal of the  
30 contents after use. Typically the kit has compartments for each useful reagent. Desirably, the reagents are provided as a dry lyophilized powder, where the reagents may be reconstituted in an aqueous medium providing appropriate concentrations of reagents for performing the assay.

35 The aforementioned constituents of the drug screening and the diagnostic assays may be used without modification or may be modified in a variety of ways. For example,

labeling may be achieved by covalently or non-covalently joining a moiety which directly or indirectly provides a detectable signal. In any of these assays, the ligand, test compound, chemokine, or antibodies thereto can be  
5 labeled either directly or indirectly. Possibilities for direct labeling include label groups: radiolabels such as  $^{125}\text{I}$ , enzymes (U.S. Pat. No. 3,645,090) such as peroxidase and alkaline phosphatase, and fluorescent labels (U.S. Pat. No. 3,940,475) capable of monitoring the change in  
10 fluorescence intensity, wavelength shift, or fluorescence polarization. Possibilities for indirect labeling include biotinylation of one constituent followed by binding to avidin coupled to one of the above label groups.

There are also numerous methods of separating bound  
15 from the free ligand, or alternatively bound from free test compound. The chemokine can be immobilized on various matrixes followed by washing. Suitable matrixes include plastic such as an ELISA plate, filters, and beads. Methods of immobilizing the chemokine to a matrix include,  
20 without limitation, direct adhesion to plastic, use of a capture antibody, chemical coupling, and biotin-avidin. The last step in this approach involves the precipitation of ligand/antibody complex by any of several methods including those utilizing, e.g., an organic solvent such as  
25 polyethylene glycol or a salt such as ammonium sulfate. Other suitable separation techniques include, without limitation, the fluorescein antibody magnetizable particle method described in Rattle, et al. (1984) Clin. Chem. 30:1457-1461, and the double antibody magnetic particle  
30 separation as described in U.S. Pat. No. 4,659,678.

Methods for linking proteins or their fragments to the various labels have been extensively reported in the literature and do not require detailed discussion here. Many of the techniques involve the use of activated  
35 carboxyl groups either through the use of carbodiimide or active esters to form peptide bonds, the formation of thioethers by reaction of a mercapto group with an

activated halogen such as chloroacetyl, or an activated olefin such as maleimide, for linkage, or the like. Fusion proteins will also find use in these applications.

Another diagnostic aspect of this invention involves  
5 use of oligonucleotide or polynucleotide sequences taken from the sequence of a chemokine. These sequences can be used as probes for detecting levels of the ligand message in samples from patients suspected of having an abnormal condition, e.g., an inflammatory or developmental problem.  
10 The preparation of both RNA and DNA nucleotide sequences, the labeling of the sequences, and the preferred size of the sequences has received ample description and discussion in the literature. Normally an oligonucleotide probe should have at least about 14 nucleotides, usually at least  
15 about 18 nucleotides, and the polynucleotide probes may be up to several kilobases. Various labels may be employed, most commonly radionuclides, particularly  $^{32}\text{P}$ . However, other techniques may also be employed, such as using biotin modified nucleotides for introduction into a  
20 polynucleotide. The biotin then serves as the site for binding to avidin or antibodies, which may be labeled with a wide variety of labels, such as radionuclides, fluorescers, enzymes, or the like. Alternatively, antibodies may be employed which can recognize specific  
25 duplexes, including DNA duplexes, RNA duplexes, DNA-RNA hybrid duplexes, or DNA-protein duplexes. The antibodies in turn may be labeled and the assay carried out where the duplex is bound to a surface, so that upon the formation of duplex on the surface, the presence of antibody bound to  
30 the duplex can be detected. The use of probes to the novel anti-sense RNA may be carried out in any conventional techniques such as nucleic acid hybridization, plus and minus screening, recombinational probing, hybrid released translation (HRT), and hybrid arrested translation (HART).  
35 This also includes amplification techniques such as polymerase chain reaction (PCR).

Diagnostic kits which also test for the qualitative or quantitative presence of other markers are also contemplated. Diagnosis or prognosis may depend on the combination of multiple indications used as markers. Thus, kits may test for combinations of markers. See, e.g., Viallet, et al. (1989) Progress in Growth Factor Res. 1:89-97.

#### X. Receptor

Having isolated a ligand binding partner of a specific interaction, methods exist for isolating the counter-partner. See, Gearing, et al EMBO J. 8:3667-4676 or McMahan, et al. (1991) EMBO J. 10:2821-2832. For example, means to label a chemokine without interfering with the binding to its receptor can be determined. For example, an affinity label can be fused to either the amino- or carboxy-terminus of the ligand. An expression library can be screened for specific binding of chemokine, e.g., by cell sorting, or other screening to detect subpopulations which express such a binding component. See, e.g., Ho, et al. (1993) Proc. Nat'l Acad. Sci. 90:11267-11271. Alternatively, a panning method may be used. See, e.g., Seed and Aruffo (1987) Proc. Nat'l. Acad. Sci. 84:3365-3369.

Protein cross-linking techniques with label can be applied to a isolate binding partners of a chemokine. This would allow identification of protein which specifically interacts with a chemokine, e.g., in a ligand-receptor like manner.

In various embodiments, new receptors designated DC CR and M/DC CR were isolated. The sequences of the human constructs and product are provided in Tables 4 and 5. Similar means for making variants and fragments, at the nucleotide level or at the protein level, and making antibodies will be available as described above, directed primarily to the chemokine embodiments. Many similar or

related uses to the ligands will be applied to the  
receptors, as specific binding reagents. In particular,  
methods will be applied to screening for specific ligands  
for each receptor. Many uses, including kits, will also be  
5 available through analogous techniques.

The broad scope of this invention is best understood with reference to the following examples, which are not intended to limit the invention to specific embodiments.

EXAMPLES

## I. General Methods

- 5        Some of the standard methods are described or  
referenced, e.g., in Maniatis, et al. (1982) Molecular  
Cloning, A Laboratory Manual, Cold Spring Harbor  
Laboratory, Cold Spring Harbor Press; Sambrook, et al.  
(1989) Molecular Cloning: A Laboratory Manual, (2d ed.),  
10 vols 1-3, CSH Press, NY; Ausubel, et al., Biology, Greene  
Publishing Associates, Brooklyn, NY; or Ausubel, et al.  
(1987 and Supplements) Current Protocols in Molecular  
Biology, Greene/Wiley, New York; Innis, et al. (eds.) (1990)  
PCR Protocols: A Guide to Methods and Applications Academic  
15 Press, N.Y. Methods for protein purification include such  
methods as ammonium sulfate precipitation, column  
chromatography, electrophoresis, centrifugation,  
crystallization, and others. See, e.g., Ausubel, et al.  
(1987 and periodic supplements); Deutscher (1990) "Guide to  
20 Protein Purification" in Methods in Enzymology, vol. 182,  
and other volumes in this series; and manufacturer's  
literature on use of protein purification products, e.g.,  
Pharmacia, Piscataway, N.J., or Bio-Rad, Richmond, CA.  
Combination with recombinant techniques allow fusion to  
25 appropriate segments, e.g., to a FLAG sequence or an  
equivalent which can be fused via a protease-removable  
sequence. See, e.g., Hochuli (1989) Chemische Industrie  
12:69-70; Hochuli (1990) "Purification of Recombinant  
Proteins with Metal Chelate Absorbent" in Setlow (ed.)  
30 Genetic Engineering, Principle and Methods 12:87-98, Plenum  
Press, N.Y.; and Crowe, et al. (1992) OIAexpress: The High  
Level Expression & Protein Purification System QUIAGEN,  
Inc., Chatsworth, CA.
- FACS analyses are described in Melamed, et al. (1990)  
35 Flow Cytometry and Sorting Wiley-Liss, Inc., New York, NY;  
Shapiro (1988) Practical Flow Cytometry Liss, New York, NY;



and Robinson, et al. (1993) Handbook of Flow Cytometry Methods Wiley-Liss, New York, NY.

## II. Isolation and characterization of chemokine cDNAs

### 5 A. TECK

The TECK was isolated from a cDNA library made from thymus cells from a RAG-1 "knockout" mouse. See, Mombaerts, et al. (1992) Cell 68:869-877. Individual cDNA clones were sequenced using standard methods, e.g., the Taq  
10 DyeDeoxy Terminator Cycle Sequencing kit (Applied Biosystems, Foster City, CA), and the TECK sequence was identified and further characterized. Computer analyses with other C-C chemokine family members revealed significant homology at the amino acid levels with other  
15 chemokines. The nucleotide sequence for mouse is provided in Table 1, encoding a polypeptide of about 144 amino acids. The signal sequence should run from 1 (met) to about 23 (ala), and removal of the signal sequence should provide one natural mature sequence beginning at 24 (gln).  
20 Additional processing may occur in a physiological system.

The sequence is notable in having a longer carboxy-terminal tail than most other CC chemokines. TECK exhibits one glycosylation site, and several AAMP, PKC, and CK2 phosphorylation sites.

### 25 B. MIP-3 $\alpha$

The MIP-3 $\alpha$  was isolated from a cDNA library made from human monocytes activated with LPS and IFN- $\gamma$ . Individual cDNA clones were sequenced using standard methods, and the MIP-3 $\alpha$  sequence was identified and further characterized.  
30 The nucleotide sequence is provided in Table 2, encoding a polypeptide of at least about 89 amino acids. The signal sequence should run from about 1 (met) to 21 (cys), and removal of the signal sequence should provide one natural sequence beginning with gly. Additional processing may  
35 occur in a physiological system.

C. MIP-3 $\beta$ 

The MIP-3 $\beta$  was isolated from a cDNA library made from human fetal lung cells. Individual cDNA clones are sequenced using standard methods, and the MIP-3 $\alpha$  sequence  
5 was identified and further characterized. The nucleotide sequence is provided in Table 3, encoding a polypeptide of about 98 amino acids. The signal sequence should run from about 1 (met) to about 21 (ser), and removal of the signal sequence should provide one mature natural sequence  
10 beginning from gly. Additional processing may occur in a physiological system.

This chemokine has been paired with a receptor designated Eb11. See Yoshida, et al. (1997) J. Biol. Chem. 13803-13809.

## 15 D. Dendritic Cell Receptor for chemokine; DC CR

The DC CR was isolated from RNA made from dendritic cells isolated from CD34<sup>+</sup> cord blood cells, isolated by standard procedure. It was also isolated from eosinophils using degenerate PCR primers of the TM2 and TM7 segments,  
20 which are often conserved among chemokine receptors. These eosinophils were isolated by taking PBLs, depletion of red blood cells by lysis, and negative selection of CD16 to remove neutrophils.

Sequencing of the PCR fragments indicated a potential  
25 novel receptor, and the fragment was used to isolate a full length clone by hybridization. Clone isolates were sequenced using standard methods, and the DC CR sequence was identified and further characterized. The nucleotide sequence is provided in Table 4, encoding a polypeptide of  
30 about 365 amino acids. The transmembrane segments, determined by homology to the IL-8 B receptor, are about: TM1 from 39 (leu) to 64 (phe); TM2 from 76 (leu) to 96 (ser); TM3 from 111 (leu) to 132 (met); TM4 from 151 (thr) to 176 (phe); TM5 from 207 (gly) to 229 (val); TM6 from 246  
35 (val) to 270 (ala); and TM7 from 291 (val) to 319 (leu). The amino terminal segment is probably an extracellular

segment, and the others would be between TM2 and TM3; and TM4 and TM5; and TM6 and TM7. The intracellular segments should then run between TM1 and TM2; TM3 and TM4, TM5 and TM6, and the carboxy terminus from the end of TM7.

- 5 Additional processing may occur in a physiological system.

The implication of chemokine receptors in retroviral infection suggest that the receptor may be critical for infection. Antibodies which block infection may be routinely screened, and developed for therapeutic uses.

- 10 E. Monocyte/Dendritic Cell Receptor for chemokine;  
M/DC CR

- The M/DC CR was isolated from a cDNA library made from human monocyte cells cultured for 2.5 to 4 h in medium containing IFN- $\gamma$  (10 ng/ml), LPS (1  $\mu$ g/ml), anti-IL-4  
15 monoclonal antibody (5  $\mu$ g/ml), and anti-IL-10 monoclonal antibody (5  $\mu$ g/ml). Individual cDNA clones were sequenced using standard methods, and the M/DC CR sequence was identified and further characterized. The nucleotide  
20 sequence is provided in Table 5, encoding a polypeptide of about 356 amino acids. The transmembrane segments, should be about as follows: TM1 from 52 (leu) to 76 (val); TM2 from 86 (asn) to 107 (ala); TM3 from 117 (ile) to 138 (val); TM4 from 157 (val) to 182 (tyr); TM5 from 211 (phe) to 233 (val); TM6 from 251 (leu) to 275 (phe); and TM7 from  
25 296 (ile) to 315 (leu). As for the DC CR, the amino terminal segment is probably an extracellular segment, and the others would be between TM2 and TM3; and TM4 and TM5; and TM6 and TM7. The intracellular segments should then run between TM1 and TM2; TM3 and TM4, TM5 and TM6, and the  
30 carboxy terminus from the end of TM7.

### III. Preparation of antibodies

- Many standard methods are available for preparation of antibodies. For example, synthetic peptides may be  
35 prepared to be used as antigen, administered to an appropriate animal, and either polyclonal or monoclonal antibodies prepared. Short peptides, e.g., less than about

10 amino acids may be repeated, while longer peptides may be used alone or conjugated to a carrier. For example, with the M/DC CR, animals were immunized with peptides corresponding to amino acid sequences from 18-44 (starting with LAP and ending with KYD; a fragment towards the amino terminus) and from 183-204 (starting with KPQ and ending with PAD; corresponding to an extracellular loop), see SEQ ID NO: 13. Highest specificity will result when the polypeptides are selected from portions which are most unique, e.g., not from conserved sequence regions. The animals may be used to collect antiserum, or may be used to generate monoclonal antibodies.

Antiserum was determined useful for ELISA, and will be evaluated for utility as immunoprecipitation or Western blot analysis. Monoclonal antibodies will also be evaluated for those same uses.

The antibodies provided will be useful as immunoaffinity reagents, as detection reagents, for immunohistochemistry, and as therapeutic reagents.

#### IV. Assays for chemotactic activity of chemokines.

Chemokine proteins are produced, e.g., in COS cells transfected with a plasmid carrying the chemokine cDNA by electroporation. See, Hara, et al. (1992) EMBO J. 10:1875-1884. Physical analytical methods may be applied, e.g., CD analysis, to compare tertiary structure to other chemokines to evaluate whether the protein has likely folded into an active conformation. After transfection, a culture supernatant is collected and subjected to bioassays. A mock control, e.g., a plasmid carrying the luciferase cDNA, is used. See, de Wet, et al. (1987) Mol. Cell. Biol. 7:725-757. A positive control, e.g., recombinant murine MIP-1 $\alpha$  from R&D Systems (Minneapolis, MN), is typically used. Likewise, antibodies may be used to block the biological activities, e.g., as a control.

Lymphocyte migration assays are performed as previously described, e.g., in Bacon, et al. (1988) Br. J.

- Pharmacol. 95:966-974. Murine Th2 T cell clones, CDC-25 (see Tony, et al. (1985) J. Exp. Med. 161:223-241) and HDK-1 (see Cherwinski, et al. (1987) J. Exp. Med. 166:1229-1244), made available from R. Coffman and A. O'Garra (DNAX, Palo Alto, CA), respectively, are used as controls.

Ca<sup>2+</sup> flux upon chemokine stimulation is measured according to the published procedure described in Bacon, et al. (1995) J. Immunol. 154:3654-3666.

- 10 Maximal numbers of migrating cells in response to MIP-1 $\alpha$  typically occur at a concentration of 10<sup>-8</sup> M, in agreement with original reports for CD4<sup>+</sup> populations of human T cells. See Schall (1993) J. Exp. Med. 177:1821-1826. A dose-response curve is determined, preferably giving a characteristic bell shaped dose-response curve.

- After stimulation with C-C chemokines, lymphocytes generally show a measurable intracellular Ca<sup>2+</sup> flux. MIP-1 $\alpha$  is capable of inducing immediate transients of calcium mobilization. Typically, the levels of chemokine used in these assays will be comparable to those used for the chemotaxis assays (1/1000 dilution of conditioned supernatants).

- 25 Retroviral infection assays have also been described, and recent description of certain chemokine receptors in retroviral infection processes may indicate that similar roles may apply to the DC CR and/or M/DC CR. See, e.g., Balter (1996) Science 272:1740 (describing evidence for chemokine receptors as coreceptors for HIV); and Deng, et al. (1996) Nature 381:661-666.

#### V. Expression analysis of chemokine/receptor genes

- RNA blot and hybridization are performed according to the standard method in Maniatis, et al. (1982) Molecular Cloning: A Laboratory Manual Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY. An appropriate fragment of a cDNA fragment is selected for use as a probe. To verify

the amount of RNA loaded in each lane, the substrate membrane is reprobed with a control cDNA, e.g., glyceraldehyde 3-phosphate dehydrogenase (G3PDH) cDNA (Clontech, Palo Alto CA).

5        Analysis of mRNA from the appropriate cell source using the probe will determine the natural size of message. It will also indicate whether different sized messages exist. The messages will be subject to analysis after isolation, e.g., by PCR or hybridization techniques.

10       Northern blot analysis may be performed on many different mRNA sources. However, in certain cases, cDNA libraries may be used to evaluate sources which are difficult to prepare. A "reverse Northern" uses cDNA inserts removed from vector, but multiplicity of bands may  
15 reflect either different sized messages, or may be artifact due to incomplete reverse transcription in the preparation of the cDNA library. In such instances, verification may be appropriate by standard Northern analysis.

20       Similarly, Southern blots may be used to evaluate species distribution of a gene. The stringency of washes of the blot will also provide information as to the extent of homology of various species counterparts.

25       Tissue distribution, and cell distribution, may be evaluated by immunohistochemistry using antibodies. Alternatively, in situ nucleic acid hybridization may also be used in such analysis.

#### A. TECK

30       The TECK was isolated from a RAG-1 "knockout" mouse. This animal is characterized by a great predominance of pro-T or pre-T cells, lacking more mature T cells after the block point of T cell receptor rearrangement. This suggests a role in very early T cell development, likely expressed by pro-T or pre-T cells, thymic stromal cells, and possibly macrophages, epithelial, and dendritic cells.  
35       This comports with the observation that tissue distribution studies have not detected significant expression in other organs or tissues. See also, Table 7.

2020/03/20




Table 7: mTECK mRNA expression in tissues and cells

cDNA libraries			northern blot		
cell type or tissue	neg	pos	cell type or tissue	neg	pos
Th2 CD4+ T cells	X		heart	X	
Th1 CD4+ T cells	X		brain	X	
Lung	X		spleen	X	
L cells	X		lung	X	
RAG-1 KO lung	X		liver	X	
RAG-1 KO heart	X		skeletal muscle	X	
RAG-1 KO brain		X (+)	kidney	X	
RAG-1 KO spleen	X		testis	X	
RAG-1 KO kidney	X		thymus		X (+++)
RAG-1 KO testis		X (+)	small intestine		X (++)
RAG-1 KO thymus		X (+++)	CD4+8- thymocytes R/A	X	
RAG-1 KO liver		X (+)	CD4+8+ thymocytes R/A	X	
CD4-8- thymocytes	X		CD4-8- thymocytes R/A	X	
A20-J B-cell lymphoma	X		B220+ splenocytes R/A	X	
BW CD4-8-3- hybridoma	X		Thy-1+ splenocytes R/A	X	
pro-T cells		X (+)	1G18LA macrophages R/A	X	
pre-T cells	X		primary thymic stroma R/A	X	
30-R bone marrow stroma	X		3D.1 thymic epithelial R/A	X	
D10 T-cell hybridoma	X		MTSC-C thymic epithelial	X	
CTLL T-cell clone	X		30.R bone marrow stroma	X	
peritoneal macrophages	X				
splenic dendritic cells	X				

Analysis of mTECK mRNA was carried out as described. + to +++ indicates the relative intensity of the signal. R/A: resting or activated.



Species analysis indicated positive signals by hybridization in human, rat, and hamster DNA. Tissue distribution analysis suggests that the gene is expressed in human small intestine, which also is a tissue which supports T cell differentiation.

The combination of the structure and distribution of this chemokine suggests a role in T cell development, which normally occurs in the thymus.

#### B. MIP-3 $\alpha$

The MIP-3 $\alpha$  was identified from a cDNA library made from human monocytes activated with LPS and IFN- $\gamma$ , in the presence of anti-IL-10. See, Rossi, et al. (1997) J. Immunology 158:1033-1036, which was published after priority dates of this filing. Message of the chemokine has also been detected in pancreatic islet cells, fetal lung, and hepatic HEPG2 cells, suggesting a physiological role in inflammation or medical conditions in such organs/tissues.

The gene is expressed in HL-60 (promyelocytic leukemia); S3 (HeLa cell); K562 (chronic myelogenous leukemia); MOLT-4 (lymphoblastic leukemia); Raji (Burkitt's lymphoma); SW480 (colorectal adenocarcinoma); A549 (lung carcinoma); and G361 (melanoma) cell lines, as determined by probing on a tissue blot from CLONTECH. Tissue expression gave a positive signal in lymph node, appendix, peripheral blood lymphocytes, fetal liver, and fetal lung, suggesting a physiological role in inflammation or medical conditions in such organs/tissues; but no detectable signal in spleen, bone marrow, brain, and kidney.

The main transcript appears to be about 1.2 kb, with two additional transcript sizes in fetal lung RNA. Among the various tissues, transcript sizes of 1.8, 2.7, and 4.2 kb were detected.

Positive signals were also detected in the following cDNA libraries: dendritic cells activated with LPS, but not when activated with GM-CSF and IL-4; monocytes treated

with LPS, IFN- $\gamma$ , and anti-IL-10, but not when treated with LPS, IFN- $\gamma$ , and IL-10; and activated PBMC.

These expression data implicate this chemokine in inflammatory responses upon cell activation. The lymph nodes, appendix, and PBL are sites where inflammatory processes take place. The MIP-3 $\alpha$  may exert its effects on monocytes and cells involved in inflammatory events. Other structural features implicate this chemokine in eosinophil and lung physiology, e.g., asthma indications. Thus, an antagonist of the chemokine, e.g., an antibody, may be important for treatment of asthmatic conditions. Also, IL-10 appears to inhibit MIP-3 $\alpha$  expression.

The human MIP-3 $\alpha$  is a ligand for the DC CR. Thus, a positive control exists for the the Ca++ flux assay for that receptor. This allows for the further screening of agonist ligands for the DC CR. Moreover, the DC CR was isolated from eosinophil cDNA, and observations have been made that eosinophils migrate to MIP-3 $\alpha$  in vitro. These suggest that the MIP-3 $\alpha$  interaction with the DC CR is important in recruitment of eosinophils, as occurs with the eotaxin ligand and the CCR3. As such, antagonists of the MIP-3 $\alpha$  interaction with the DC CR will likely be useful in inhibiting eosinophilia, particularly in the lung, or lung inflammation. These may accompany asthmatic or other pulmonary conditions.

Antagonists to MIP-3 $\alpha$  may be made either with antibodies, or other binding compositions which inhibit receptor interaction. The antibodies may be to the ligand, MIP-3 $\alpha$  itself, or to the binding portions of the receptor, DC CR. Muteins of the chemokine may block receptor interaction, and with a positive control, chemokine muteins may be screened for variations which compete with the wild type chemokine at various concentrations. See, e.g., Kenakin (1987) Pharmacological Analysis of Drug-Receptor Interaction Raven Press, NY; Arunlakshana and Schild (1959) Br. J. Pharmacol. 14:48-58; Black (1989) Science 245:486-493; Zurawski, et al. (1986) J. Immunol. 137:3354-3360;

Zurawski and Zurawski (1988) EMBO J. 7:1061-1069; Zurawski and Zurawski (1992) EMBO J. 11:3905-3910; Imler and Zurawski (1992) J. Biol. Chem. 267:13185-13190.

C. MIP-3 $\beta$

- 5       The MIP-3 $\beta$  was identified in a cDNA library made from human monocytes activated with LPS and IFN- $\gamma$ , in the presence of anti-IL-10. Its distribution in other cells and tissues has not been fully determined.

D. Dendritic Cell Receptor for chemokine; DC CR

- 10       The DC CR was isolated from a cDNA library made from a dendritic cell cDNA library. It appears to be expressed in certain T cells, spleen cell subsets, NK cells, and other cell populations enriched in dendritic cells, including CD1a<sup>+</sup>, CD14<sup>+</sup>, and CD1Aa<sup>+</sup> cells. It did not give a  
15 detectable signal in TF1, Jurkat, MRC5, JY, or U937 cells.

- Being found on dendritic cells, its ligand, including the MIP-3 $\alpha$ , may be important in attracting appropriate cells for the initiation of an immune response. MIP-3 $\alpha$  has been shown to be a very potent chemoattractant for  
20 dendritic cells. Significant roles of the ligand and receptor in pulmonary physiology are suggested, both from the distribution of the receptor and ligand. The receptor may be also present in other cells important in such responses.

- 25       E. Monocyte/Dendritic Cell Receptor for chemokine; M/DC CR

- The M/DC CR was isolated from a cDNA library made from primary monocyte cells activated with LPS and IFN- $\gamma$  but subtracted with known high abundance genes from those  
30 cells. The abundance of this gene is probably less than about 1% of message from those cells.

- Tissue expression gave a positive signal in spleen, PBL, lung, placenta, and small intestine; but no detectable signal in brain, liver, kidney, and muscle. This  
35 distribution suggests a hematopoietic role.

There appears to be one main transcript, but the existence of additional or alternatively spliced messages has not been eliminated.

Positive signals were also detected in the following  
5 cDNA libraries: monocytes and dendritic cells; but signals  
were not detectable in CD8<sup>+</sup> T cells, or in either resting  
or activated splenocytes, gamma-delta T cells, NK cells, or  
B cells. Immunohistochemistry will be performed to confirm  
absence in the T cell and B cell compartments and to check  
10 in tonsil, particularly in view of location in spleen and  
placenta. The relatively restricted distribution on  
monocytes and dendritic cells leads both to its  
designation, and suggests a functional role in those cell  
types, which are important in the initiation of immune  
15 responses through their ability to process and present  
antigen to T cells.

#### VI. Specific Characterization of TECK

A novel CC chemokine was identified in the thymus of  
20 mouse and human and was designated TECK as Thymus Expressed  
ChemoKine. TECK has weak homology with other CC chemokines  
and maps to mouse chromosome 8. Besides the thymus, mRNA  
encoding TECK was detected at substantial levels in the  
small intestine and at low levels in the liver. The source  
25 of TECK in the thymus was determined to be thymic dendritic  
cells, while in contrast bone marrow-derived dendritic  
cells do not express TECK. The murine TECK recombinant  
protein showed chemotactic activity for activated  
macrophages, dendritic cells and thymocytes. We conclude  
30 that TECK represents a novel thymic dendritic cell-specific  
CC chemokine which is possibly involved in T-cell  
development.

Chemokines belong to a family of small peptides (6-15  
kDa) whose best described biological function is to control  
35 the migration of certain leukocyte populations to localized  
sites of inflammation. Baggiolini, et al. (1994) Adv. in  
Immun. 55:97-179; Schall and Bacon (1994) Curr Opin Immun

6:865-873; Hedrick and Zlotnik (1996) Curr. Opin. Immunol.  
8:343-347. In the last few years many new members of the  
chemokine super family have been characterized.  
Initially, new chemokines were identified through their  
5 chemotactic effects on leukocytes (Baggiolini et al.  
(1994); Schall and Bacon (1994)) and were isolated mainly  
from blood leukocytes or cell lines. More recently,  
approaches based on the selective cloning of secreted  
molecules by signal sequence trap (Tashiro, et al. (1993)  
10 Science 261:600-603; Imai, et al. (1996) J. Biol. Chem.  
271:21514-21521) or on the exploitation of public and  
private databases of expressed sequence tags (EST) through  
bioinformatics (Hieshima, et al. (1997) J. Biol. Chem.  
272:5846-5853; Patel, et al. (1997) J. Exp. Med. 185:1163-  
15 1172; and Rossi, et al. (1997) J. Immunol. 158:1033-1036),  
have allowed the rapid identification of novel chemokines  
based on sequence and structural homologies. These  
approaches take advantage of the fact that most of the  
chemokines are secreted factors whose protein sequence  
20 contain four conserved cysteines (Schall (1994) "The  
Chemokines" pp. 419-460 in Thomson (eds.) The Cytokine  
Handbook, Academic Press, New York. The CXC or  $\alpha$  chemokine  
family has the two first amino-terminal cysteines separated  
by a non-conserved amino acid. In the CC or  $\beta$  chemokine  
25 family, these two cysteines are consecutive. A third type  
of chemokine, the C or  $\gamma$  family, is represented by  
lymphotactin, which conserves two cysteines (1 and 3)  
instead of the original four (Kelner, et al. (1994) Science  
266:1395-1399). Finally, a recently identified chemokine  
30 with three amino acids separating the first two cysteines  
defines a fourth CX<sub>3</sub>C family (Bazan, et al. (1997) Nature  
385:640-644).

Interestingly, some of the new chemokines discovered  
show a relatively restricted pattern of expression (Imai et  
35 al. (1996); Hieshima et al. (1997)). It is tempting to  
suggest that these new approaches may lead to the discovery  
of tissue- or cell-specific chemokines. In addition, new

biological evidence for important new roles of chemokines in hemopoiesis (Cook (1996) J. Leukoc. Biol. 59:61-66; and Nagasawa, et al. (1996) Nature 382:635-638) and the control of viral infections including HIV (Cocchi, et al. (1995) Science 270:1811-1815; and Cook, et al. (1995) Science 269:1583-1585). Thus, the molecular cloning of novel chemokines through DNA-based strategies may uncover novel proteins belonging to the chemokine super family but whose physiological role goes beyond the control of inflammation.

10 In an attempt to identify novel genes involved in T-cell development, we analyzed a cDNA library from the thymus of Recombinase Activation Gene-1 (RAG-1) deficient mice. We identified a novel CC chemokine designated TECK for Thymus Expressed ChemoKine, based on sequence homology with other known chemokines. We subsequently isolated the human homologue of TECK. The pattern of expression of TECK mRNA is highly restricted to the thymus and small intestine in both human and mouse. Moreover, in the mouse thymus, TECK protein is produced by dendritic cells while splenic dendritic cells do not express TECK mRNA. Recombinant TECK showed chemotactic activity on thymocytes, macrophages, THP-1 cells and dendritic cells, while it was inactive on peripheral lymphocytes and neutrophils. The restricted pattern of expression of TECK together with its biological properties suggest a role for this novel dendritic cell-specific chemokine in T-cell development.

#### A. Cloning and structural analysis of mouse TECK

A directional cDNA library was made from RAG-1 deficient mouse thymus and analyzed by random sequencing. One of the clones contained an open reading frame with significant homology to previously described CC chemokines. The full-length cDNA contains 1037 bp including an open reading frame of 426 bp encoding a protein of 142 amino acids and will be identified in this report as mTECK (see Table 1). In the 3' untranslated region, there is one unique polyadenylation signal consistent with the single mRNA species observed in northern blots. The mTECK cDNA

does not contain any ATTTA transcript destabilization motif (Shaw and Kamrn (1986) Cell 46:659-667). The comparison of the amino acid sequence of mTECK with previously described murine CC chemokines demonstrates the conservation of the four cysteines present in all these chemokines. However, mTECK shows few additional identities with these proteins.

B. Cloning and molecular characterization of human TECK

To investigate the possible existence of a gene homologous to mTECK in other mammalian species, a Southern blot with genomic DNA from various species was hybridized with the mTECK cDNA probe. Under high stringency conditions, hybridizing bands were detected in mouse, rat, hamster and human genomic DNAs. Interestingly, a single band was detected in human, suggesting that a single gene encodes for TECK in this species. The multiple bands present in mouse, rat and hamster could be the result of a internal EcoRI site within the TECK gene. Alternatively, the TECK gene may have been duplicated in these species.

In order to clone the human homologue of mTECK, a blot of cDNAs from a panel of human cDNA libraries was hybridized with the mTECK cDNA probe. A signal was observed in a fetal small intestine cDNA library. Screening of this library with the mTECK probe allowed the isolation of several identical clones of 1012 bp with an open reading frame of 453 bp encoding a protein of 151 amino acids. This protein had a much higher degree of homology at the nucleic acid level (71% nucleic acid identity for the open reading frame and 49.3% amino acid identity) to mTECK than to other known CC chemokines and was thus designated as hTECK.

#### C. Chromosomal location of mTECK

It has been shown that the genes encoding for most chemokines are clustered in the genome. The genes encoding CC chemokines cluster on mouse chromosome 11 and human chromosome 17q11-12. Schall (1994); and Hedrick and Zlotnik (1996). The chromosomal location of mTECK (designated gene symbol Teck) was determined by inter

specific backcross analysis between ([C57Bl/6J X M.  
spretus]F1 X C57Bl/6J) mice. Jenkins, et al. (1982) J.  
Virol. 43:26-36. The mapping results indicated that the  
Teck locus is located on the proximal part of mouse  
5 chromosome 8. Although the chromosomal location of the  
human Teck locus could not be determined, this region of  
mouse chromosome 8 is syntenic to the human 19p13.3 and  
13q34 regions. However, the Teck locus is also very close  
to a region syntenic to human chromosome 4. The closest  
10 known gene, Insr, encodes the insulin receptor and the  
genetic distance between Teck and Insr was estimated at  $0.9 \pm 0.9$  cM. We have compared our inter specific map of  
chromosome 8 with a composite mouse linkage map that  
reports the map location of many uncloned mouse mutations  
15 (Mouse Genome Database, The Jackson Laboratory, Bar Harbor,  
ME). Teck resides in a region of the composite map that  
lacks mouse mutations with a phenotype that might be  
expected for an alteration in this locus.

D. Analysis of mTECK and hTECK mRNA distribution in cells  
20 and tissues

An analysis of the distribution of mTECK mRNA in  
tissues and cells by northern blotting or by Southern  
blotting of mouse cDNA libraries revealed that mTECK was  
expressed at significant levels only in the thymus and to a  
25 lesser extent in small intestine (Table 7). Weak  
expression of mTECK mRNA was observed in brain, testis, and  
liver RAG-1-/- cDNA libraries. Interestingly, mTECK mRNA  
was detected in a cDNA library of activated pro-T cells.  
Pro-T cells represent an early stage of intra-thymic T-cell  
30 progenitors, not fully committed to the T-cell lineage  
since they can give rise to NK and dendritic cells. Moore  
and Zlotnik (1995) Blood 86:1850-1860; Wu, et al. (1996).  
J. Exp. Med. 184:903-911). In contrast, mTECK mRNA was  
undetectable in resting or activated thymocytes, peripheral  
35 T or B cells, macrophages, PBLs, splenic dendritic cells  
and in all other tissues tested, with the exception of  
spleens recovered from mice injected with LPS (Table 7).



Interestingly, mTECK mRNA was detected by PCR in fetal thymuses of day 14 of gestation, indicating that mTECK is expressed in the thymus at the earliest stages of T-cell development.

5        The distribution of hTECK mRNA was similarly analyzed. As with mouse, hTECK mRNA expression was highly restricted to the thymus and small intestine. Weak expression was also detected in inflamed tonsil and fetal spleen, but at much lower levels than that observed in the thymus since  
10 this particular blot was exposed for a long time.

Importantly, hTECK mRNA was absent from a series of cDNA libraries from dendritic cells derived in vitro from bone marrow CD34+ progenitors cells or peripheral blood monocytes. In addition, hTECK mRNA was also absent from  
15 libraries of monocyte-derived dendritic cells stimulated with LPS or a combination of TNF- $\alpha$ , IL-1 $\alpha$  and monocyte supernatant for 4 and 16 hours. Collectively, these data indicate that TECK mRNA is specifically expressed at high levels in thymus and small intestine in vivo.

20 E. Identification of mTECK-producing cells in vivo

The abundance of mTECK mRNA expression in RAG-1 deficient thymus and its absence in thymic T cells suggested that mTECK was expressed by a thymic stromal component in normal mice. We performed in situ mRNA  
25 hybridization with sense or antisense mTECK probes generated by PCR. Thymic sections hybridized with the sense probe (negative control) demonstrated no specific staining while sections hybridized with the anti-sense probe at the same concentration showed specific staining in  
30 the thymic medulla. At higher magnification, positive cells appeared to have a non-lymphoid morphology with processes surrounding lymphoid cells. This experiment indicated that, in vivo, mTECK mRNA is expressed by a non-lymphoid component of the medullary stroma, possibly  
35 dendritic cells.

The thymic stroma is mainly composed of epithelial cells, macrophages, dendritic cells and fibroblasts,

together with a network of vascular and nervous tissue.  
Boyd, et al. (1993) Immunol. Today 14:445-459. Since we  
previously failed to detect mTECK mRNA expression in thymic  
epithelial or macrophage cell lines with or without  
5 activation with IFN- $\gamma$  (Table 7), we sorted thymic dendritic  
cells based on their high expression of MHC class II and  
CD11c (N-418 antibody). Analysis of mTECK expression by  
RT-PCR revealed that freshly isolated MHC class II+ CD11c+  
thymic dendritic cells expressed mTECK mRNA while the MHC  
10 class II+ CD11c- subset was negative. In contrast, mTECK  
mRNA was undetectable in a cDNA library made from freshly  
isolated splenic dendritic cells (Table 7).

We then performed immunostaining of thymic sections  
and purified thymic dendritic cells with a polyclonal  
15 antibody raised against a decapeptide corresponding to the  
C-terminus of mTECK. This polyclonal antibody reacted with  
recombinant mTECK in ELISA and western blot while normal  
rabbit serum was negative. In thymic sections, the  
polyclonal anti-peptide antibody reacted with a stromal  
20 component of the thymic medulla consistent with the in situ  
hybridization data while staining with normal rabbit serum  
was negative. Interestingly, the antibody also reacted  
weakly with some endothelial cells, raising the possibility  
that mTECK can be produced by the thymic endothelium.  
25 Finally, the anti-mTECK polyclonal antibody stained sorted  
thymic dendritic cells, while the control serum was  
negative. High magnification clearly showed intra-cellular  
staining of cells with characteristic dendritic morphology.  
Taken together, these results indicate that thymic  
30 dendritic cells and possibly thymic endothelial cells are  
producing TECK in vivo.

#### F. Chemotactic activities of mTECK protein

To evaluate the biologic properties of mTECK, a  
recombinant protein with a N-terminal FLAG peptide was  
35 obtained in a bacterial expression system. In some  
experiments, a recombinant mTECK protein with a C-terminal  
FLAG was used and similar results were obtained.

Interestingly, mTECK induced the migration of mouse thymocytes (Figure 1A). The optimal response was obtained with a dose of 10 ng/ml TECK. Cell migration was determined to be chemotaxis and not chemokinesis through the checkerboard analysis. Furthermore, it is established that chemokines bind to specific receptors that are coupled through heterotrimeric G proteins to intra-cellular signal-transducing pathways. Murphy (1994) Annu. Rev. Immunol. 12:593-633. To determine whether the chemotaxis of thymocytes involved a G protein-coupled receptor, cells were incubated prior to the assay with 10 ng/ml pertussis toxin which ADP-ribosylates G $\alpha$ i-proteins. Katz, et al. (1992) Nature 360:686-689. This pre-treatment completely abrogated the chemotactic response of thymocytes to mTECK (Figure 1A).

The recombinant mTECK protein also induced the migration of human monocytic THP-1 cells activated for 16 hours with IFN- $\gamma$  (Figure 1B), while it was not significantly active on resting THP-1 cells. This experiment showed that mTECK is active on human cells. In addition, mTECK induced activated mouse peritoneal macrophages to migrate as well as highly purified mouse splenic dendritic cells (Figure 1B). In all these experiments, the optimal dose of mTECK was 10 ng/ml. In contrast, no chemotaxis was observed with bone marrow cells, purified neutrophils, splenic B cells, splenic T cells or IL-2 activated RAG-1 deficient mouse splenocytes lacking mature T and B lymphocytes (Mombaerts, et al. (1992) Cell 68:869-877) and therefore enriched in NK cells. These data are consistent with the absence of in vivo accumulation of neutrophils, monocytes or lymphocytes 2 and 5 h following an intra-peritoneal injection of 10  $\mu$ g mTECK. Collectively, these data indicate that TECK is a chemotactic factor for thymocytes, macrophages and dendritic cells.

G. TECK, a distant member of the CC chemokine family

In this report, we describe the molecular isolation and characterization of TECK, a novel mouse and human CC

chemokine. Analysis of its predicted amino acid sequence showed that TECK is distantly related to previously described CC chemokines. Conservation of particular amino acids among most CC chemokines may be related to their functional importance. Beall, et al. (1992) J. Biol. Chem. 267:3455-3459; and Lusti-Narasimhan, et al. (1995) J. Biol. Chem. 270:2716-2721. In particular, a tyrosine residue between the second and third cysteines has been shown to be critical for monocyte chemotaxis (in position 50) (Beall et al. (1992)). While TECK does not have a tyrosine at this particular position, it has one in position 52 that may have the same function, since TECK is chemotactic for activated monocytes. In addition to these differences in the primary structure, the gene encoding TECK maps on chromosome 8 in the mouse, unlike most other CC chemokines which are clustered on chromosome 11. This is not the first report of an unusual chromosomal location for a CC chemokine. We have cloned the human CC chemokine MIP-3 $\beta$  and showed that its gene was on chromosome 9 rather than 17 (Rossi, et al. (1997)), and the gene encoding the novel human CC chemokine MIP-3 $\alpha$ /LARC (Rossi, et al. (1997)) has been mapped on chromosome 2 (Hieshima, et al. (1997)). It is likely that the CC chemokines on chromosome 11 in the mouse and 17 in human have been generated through gene duplication of a primordial chemokine. Our results suggest that TECK may have been generated at an earlier stage during evolution. In this regard, the TECK gene may have evolved to ensure functions similar to other CC chemokines with a distant primary structure but through similar receptor(s) as dictated by its secondary and tertiary structures. Alternatively, the receptor(s) and physiological role of TECK may be unique among chemokines.

H. TECK expression and function is associated with T-cell development

We observed that TECK was strongly expressed in the thymus which is the primary lymphoid organ where T-cell development takes place. Recently, another CC chemokine

highly expressed in the thymus, TECK, has been identified. Imai, et al. (1996). However, TECK is also expressed in lung and colon as well as activated PBMC (Imai, et al. (1996)) while TECK was absent from these tissues. Besides  
5 the thymus, numerous reports indicate that T cell development can occur in the small intestine (Poussier and Julius (1994) Annu. Rev. Immunol. 12:521-553) where TECK is also expressed. Interestingly, the liver has also been suggested to support T-cell development to some extent  
10 (Abo, et al. (1994) Int. Rev. Immunol. 11:61-102) and we observed a low TECK expression in a liver cDNA library. These data show that TECK expression correlates with organs that support T-cell development.

While many molecular and cellular aspects of T-cell  
15 differentiation are well documented, the precise role of chemokines in T-cell development is still unknown. Recently, it has been shown that the bone marrow stroma-derived CXC chemokine SDF-1 is important for B lymphopoiesis and myelopoiesis since SDF-1 -/- mice are  
20 impaired for these functions (Nagasawa, et al. (1996)). Similarly, it is likely that chemokines act at different steps of T-cell differentiation. Chemokines, together with the expression of appropriate adhesion molecules, may dictate the migration of uncommitted progenitors from the  
25 bone marrow to other anatomic locations. Indeed SDF-1 is chemoattractant for human CD34+ progenitor cells. Aiuti, et al. (1997) J. Exp. Med. 185:111-120. The observation that TECK is chemoattractant for thymocytes but not for mature peripheral T cells suggests that TECK could attract  
30 T-cell progenitors to the thymus. Such populations are very difficult to isolate in sufficient numbers to conduct in vitro chemotaxis experiments, but we are currently designing new strategies to address this important question. In addition, we have not found significant  
35 chemotactic activity of TECK on bone marrow cells. SDF-1 was shown to be much less potent on CD34+ progenitors from the peripheral blood than those from the bone marrow.

Aiuti, et al. (1997). It is possible that the sensitivity of progenitor cells to TECK would increase as these cells leave the bone marrow to colonize lymphoid organs.

Importantly, intra-thymic maturation is also characterized  
5 by a directional migration from the subcapsular region which contains the earliest progenitors to the cortex and finally to the medulla where thymocytes finish their maturation (Boyd, et al. (1993)). It is possible that the  
10 secretion of TECK by medullary dendritic cells may play a role in this directional migration. Yet another possibility is that TECK may play a role in the organization and development of the thymic stroma.

We also showed that TECK is chemotactic for activated  
15 macrophages and dendritic cells. These two cell types also play important roles in T-cell development. Through a complex screening process involving positive and negative selection events most of the antigenic specificities randomly generated in the thymus will be eliminated by  
20 programmed cell death (Janeway (1994) Immunity 1:3-6). The efficient scavenging of dead thymocytes is probably mediated, at least partially, by thymic macrophages and thus TECK could play an important role through its action on activated macrophages. Further along, T-cells with a high affinity for self-antigens and thus potentially  
25 harmful are eliminated through negative selection (Janeway (1994)). It is believed that thymic dendritic cells are primarily responsible for the negative selection of thymocytes, therefore playing a major role in the establishment of tolerance. Inaba, et al. (1991) J. Exp.  
30 Med. 173:549-559. An efficient mechanism of central tolerance should eliminate T cells potentially reactive against auto-antigens which are not expressed in the thymus, such as organ specific auto-antigens. Several known chemokines induce the migration of dendritic cells  
35 and could therefore contribute to their recruitment during peripheral immune responses. Sozzani, et al. (1995) J. Immunol. 155:3292-3295; and Xu, et al. (1996) J. Leukoc.

Biol. 60:365-371. Similarly, dendritic cells presenting organ-specific or other antigens could be recruited to the thymus or the small intestine and induce negative selection of T cells specific for these antigens. It is possible that thymus- and small intestine-specific chemokines active on dendritic cells such as TECK could play an important role in the establishment of tolerance. Thus, TECK could potentially interact at several important steps of T-cell development. Future experiments will aim to define the precise role of TECK in T-cell development and other physiological processes through the use of genetically modified mice.

I. TECK is specifically expressed by thymic dendritic cells

Dendritic cells represent an heterogeneous cell population derived from bone marrow progenitors. They are present in non-lymphoid organs as immature dendritic cells (such as Langerhans cells in the skin) where they display a high ability for antigen capture. Cella, et al. (1997) Curr. Opin. Immunol. 9:10-16. Subsequent to antigen challenge, they will migrate to secondary lymphoid organs and will acquire a high capacity to present processed antigens to naive T-cells to initiate a specific immune response (Cella, et al. (1997)). It has been shown that dendritic cells can derive from CD34+ progenitors cultured in the presence of GM-CSF and TNF- $\alpha$  (Caux, et al. (1992) Nature 360:258-261; and Caux, et al. (1996) J. Exp. Med. 184:695-706) or from monocytes in the presence of GM-CSF and IL-4 (Sallusto and Lanzavecchia (1994) J. Exp. Med. 179:1109-1118). Interestingly, there is also evidence for a lymphoid dendritic cell precursor in thymus and bone marrow which is able to derive both lymphocytes and dendritic cells in the absence of GM-CSF. Ardavin, et al. (1993) Nature 362:761-763; Galy, et al. (1995) Immunity 3:459-473; Marquez, et al. (1995) J. Exp. Med. 181:475-483; and Wu, et al. (1996). These lymphoid-derived dendritic cells may have different functional properties such as a

negative regulation of T-cell responses since they express FasL in the mouse. Suss and Shortman (1996) J. Exp. Med. 183:1789-1796. We found that TECK was expressed at high levels in mouse thymic dendritic cells but was absent in cDNA libraries from mouse splenic dendritic cells or from human dendritic cells generated in vitro from CD34+ precursors or monocytes. Interestingly, mTECK mRNA was present at a low level in a population of early thymocyte progenitors still able to derive dendritic cells (Wu, et al. (1996). Thus, it would be tempting to suggest that TECK could be a specific marker of lymphoid-derived dendritic cells. However, we observed that TECK was absent from splenic dendritic cells that likely contain lymphoid-derived dendritic cells. The expression of TECK mRNA appeared in the spleen of mice injected with LPS would suggest that peripheral dendritic cells may express TECK upon activation, but we found that TECK was not expressed in cDNA libraries of bone-marrow derived dendritic cells activated with LPS, PMA and ionomycin or IL-1 $\alpha$  and TNF- $\alpha$ . It is possible that the normal expression of TECK is specific for lymphoid-derived dendritic cells or, alternatively, that it is upregulated by very specific stimuli present in the thymic and intestinal micro-environment under physiological conditions. Consistent with the latter hypothesis is our observation of specific staining of thymic endothelial cells with anti-TECK antibody since we have not been able to find TECK expression in human HUVEC endothelial cells by northern blot analysis, without activation or following a 16 hour-activation with various combinations of IL-1, TNF- $\alpha$ , IL-4, IL-7 and oncostatin while some of these stimuli induce the expression of other CC chemokines in endothelial cells. Rollins and Pober (1991) Am. J. Pathol. 138:1315-1319; Marfaing-Koka, et al. (1995) J. Immunol. 154:1870-1878; Garcia-Zepeda, et al. (1996) J. Immunol. 157:5613-5626; and Garcia-Zepeda, et al. (1996) Nat. Med. 4:449-456. Taken together, our data strongly suggest that TECK is a novel



chemokine specifically expressed by activated lymphoid-derived dendritic cells.

Through their function of antigen presentation, dendritic cells play major roles in the establishment of tolerance and in the initiation of an antigen-specific immune response. The use of purified dendritic cells has been recently proposed in different therapeutic protocols (Cella, et al. (1997)). The discovery of factors with a regulated expression in dendritic cells such as the novel CC chemokine TECK will certainly improve our knowledge of the biology of dendritic cells and lead to the design of relevant in vivo applications.

#### J. Mice and in vivo experimental procedures

Four to eight week-old and time-pregnant BALB/c mice were purchased from Simonsen Laboratories (Gilroy, CA). RAG-1-deficient mice (Mombaerts, et al. (1992)) were purchased from The Jackson Laboratory (Bar Harbor, ME). To analyze TECK expression after in vivo activation, various organs were recovered from pools of 2 mice 3 hours after intravenous LPS injection (50 µg LPS in 200 µl PBS or 200 µl PBS for controls).

#### K. Cell purification, culture and stimulation.

THP-1 cells (TIB-202 from the American Type Culture Collection, Rockville, MD) were cultured in complete medium which consisted in RPMI 1640 medium (JRH BioSciences, Lenexa, KS) supplemented with 10% FCS, 200 mM L-glutamin,  $5 \times 10^{-5}$  M mercaptoethanol, MEM amino-acids and vitamins, sodium bicarbonate, penicillin, streptomycin (all from Sigma, ST. Louis, MO), and gentamycin (Boehringer, Indianapolis, IN). To obtain activated mouse macrophages, 10 ml of cold PBS were injected into the peritoneum and the collected cells allowed to adhere to plastic for 24 h in complete medium. The adherent fraction, mostly macrophages, was then collected. To obtain splenic dendritic cells, a splenocyte cell suspension was prepared in RPMI 1640 Dutch modified medium (Life Technologies,

Paisley, Scotland) as described previously in, e.g.,  
Macatonia, et al. (1987) J. Exp. Med. 166:1654-1667.

Splenocytes were incubated at 37° C for 16 h and the cell  
suspension was collected and laid over Metrizamide (Nycomed  
5 Pharma, Oslo, Norway). After centrifugation for 10 min. at  
1700 x g, the low interface was collected and stained with  
anti-Mac-1 (Pharmingen, San Diego, CA) and the anti-CD11c  
N-418 antibodies (Macatonia, et al. (1993) J. Immunol.  
150:3755-3765). Splenic dendritic cells were sorted by  
10 flow cytometry on a FACStar plus cell sorter (Becton  
Dickinson, Mountain View, CA) to a purity greater than 98%  
upon reanalysis in all the experiments included in this  
report. To obtain thymic dendritic cells, thymuses were  
cut in small fragments and resuspended in 10 ml of RPMI-  
15 1640 +10% FCS containing 1 mg/ml collagenase and 0.02 mg/ml  
DNase I (both from SIGMA) and digested with continuous  
agitation at room temperature for 30 min. (Shortman, et  
al. (1995) Adv. Exp. Med. Biol. 378:21-29). One ml of 0.1M  
EDTA pH 7.2 was added for an additional 5 min. Cells were  
20 then washed in complete medium, resuspended in complete  
medium and overlaid onto Metrizamide. The thymic dendritic  
cell-enriched preparation was then stained with anti-IAD  
and N-418 antibodies and the dendritic cells sorted by flow  
cytometry

## 25 L. Molecular cloning of mouse and human TECK

The cDNA encoding mouse TECK was obtained by random  
sequencing of a RAG-1 KO mouse thymic directional cDNA  
library. Briefly, mRNA was extracted using RNazol™ B  
(Tel-Test, Friendswood, TX) and then oligotex-dT mRNA kit  
30 (Quiagen, Chatsworth, CA) following the manufacturer's  
instruction. A directional cDNA library was prepared using  
the Superscript™ Plasmid System (Gibco-BRL, Grand Island,  
NY) and cloned into the pME18s plasmid vector. Sequencing  
was done using the TaQ DyeDeoxy Terminator Cycle Sequencing  
35 kit (Applied Biosystems, Foster City, CA). To determine  
whether TECK was present in other mammals including human,  
a Southern blot containing EcoRI digested genomic DNA from

different species (Bios Laboratories, New Haven, CT) was hybridized with the full-length mouse TECK cDNA.

The cDNA encoding human TECK was found by screening of a small intestine cDNA library using the full-length mouse  
5 TECK cDNA as a probe following standard procedures.

M. Northern blot analysis of RNA and Southern blot analysis of cDNA libraries

All RNA's were isolated from tissues or cells using RNAzol™ B (Tel-Test) and analyzed after electrophoresis in  
10 a 1% formaldehyde-agarose gel (10 µg/lane). RNA's were then blotted onto a Hybond-N+ nylon membrane (Amersham, Arlington Heights, IL). Some northern blots of mRNA were bought from Clontech (Palo Alto, CA). To analyze the expression of TECK in cDNA libraries (obtained from T.  
15 MacClanahan, DNAX), 10 µg of DNA were digested with the appropriate restriction enzymes to release their inserts and analyze by Southern blotting onto nylon membranes. Northern blots and blots of cDNA libraries were hybridized for 16 hours at 65°C with a <sup>32</sup>P-labeled probe consisting in  
20 the full-length cDNA encoding for mouse or human TECK and then washed and exposed, according to standard protocols.

N. Inter specific mouse backcross mapping

Inter specific backcross progeny were generated by mating (C57Bl/6J x M. spretus) F1 females and C57Bl/6J  
25 males as described, e.g., in Copeland and Jenkins (1991) Trends Genet. 7:113-118. A total of 205 N<sub>2</sub> mice were used to map the Teck locus. DNA isolation, restriction enzyme digestion, agarose gel electrophoresis, Southern blot transfer and hybridization with the full-length mTECK cDNA  
30 probe were performed as described, e.g., in Jenkins, et al. (1982). Fragments of 7.5, 6.9, and 2.5 kb were detected in HincII digested C57Bl/6J DNA and fragments of 8.8 and 5.4 kb were detected in HincII digested M. spretus DNA. The presence or absence of the 8.8 and 5.4 kb HincII M.  
35 spretus-specific fragments, which cosegregated, was followed in backcross mice. A description of the probes and RFLPs for two of the loci linked to Teck including Insr

has been reported previously, e.g., in Ceci, et al. (1990) Genomics 6:72-79. Recombination distances were calculated as described (Green (1981) "Linkage, recombination and mapping" pp. 77-113 in Genetics and Probability in Animal Breeding Experiments, Oxford University Press, New York) using the computer program SPRETUS MADNESS.

O. Measurement of TECK mRNA expression by RT-PCR

RNA's from sorted thymic dendritic cells or fetal thymuses were prepared with the RNeasy total RNA kit (Quiagen, Chatsworth, CA), following the manufacturer's instructions. First strand cDNAs were generated by reverse transcription with a random hexamer in a 10 µl reaction and 1 µl of this reaction was used as a template for PCR. TECK expression was compared to the expression of hypoxanthine-guanine phosphoribosyl transferase (HPRT). Primer sequences were as follows: TECK: 5' primer, 5'CCTTCAGGTATCTGGAGAGGAGATC3' (nucleotides 58-72 of SEQ ID NO: 1) and 3' primer, 5'CACGCTTGTTGTTGGGGTTC3' (complement of nucleotides 447-468 of SEQ ID NO: 1), HPRT: 5' primer, 5'GTAATGATCAGTCAACGGGGGAC3' (SEQ ID NO: 17) and 3' primer, 5'CCAGCAAGCTTGCAACCTTAACCA3' (SEQ ID NO: 18). Samples were submitted to 25 cycles of amplification, each composed of 94° C for 1 min., 57° C for 30 s and 72° C for 2 min. PCR products were then separated by electrophoresis in 2% agarose gels and stained with ethidium bromide.

P. In Situ Hybridization

Biotin-14-CTP labeled sense and antisense riboprobes were generated using a non radioactive RNA labeling system (Gibco, Gaithersburg, MD) and the plasmid PCRII. (InVitrogen, Carlsbad, CA) containing a 400 base pair TECK cDNA fragment inserted by PCR and TA cloning (InVitrogen). Paraffin-embedded tissues were cut in 3-5 µm sections, mounted on slides, baked at 60° C for one hour, deparaffinized in xylene (Fisher Scientific, Pittsburgh, PA) and immersed in 100% ethanol. Sections were then incubated for 10 min at 37° C in proteinase K solution (40 mg/ml) (Gibco) in PBS and rinsed for 2 min in PBS at room

temperature before being refixed in 10% formalin (Fisher Scientific, Pittsburgh, PA) in PBS for 1 min. Next, the sections were dehydrated through graded solutions of ethanol and air dried. Hybridization was carried out using the Gibco in situ hybridization and detection system kit. Vanadyl ribonucleoside complex (Gibco) was added to the hybridization solution (39 mM final). A 0.1 µg/ml concentration of each probe was used during an 18 h hybridization at 42° C. Post-hybridization washes used room temperature 0.2X SSC. Following detection and substrate visualization, the slides were counterstained with 1% nuclear red stain (Sigma, St. Louis, MO).

#### Q. Immunohistochemistry

A polyclonal antibody specific of a synthetic decapeptide identical to the C-terminus part of murine TECK (Figure 1) was prepared in rabbits by Research Genetics (Huntsville, AL). Normal rabbit serum from a pool of 50 different animals (Research Genetics) was used as a negative control. To study TECK protein expression in the mouse thymus, 6 µm thick cryostat sections were thaw mounted on organosilicone subbed slides (American Histology Reagent Co., Stockton, CA.) and fixed in 3% formalin (Fisher Scientific, Springfield, NJ) in Hank's Balanced Salt Solution with 0.01M HEPES (HBSS-HEPES), pH 7.4, for 15 min at room temperature. The sections were sequentially blocked for endogenous biotin binding using the Vector blocking kit (Vector Laboratories, Burlingame, CA) and for endogenous peroxidase activity with a 1% hydrogen peroxide, 0.2M sodium azide solution, in HBSS-HEPES with 0.1% saponin (staining buffer). Non-specific antibody binding sites were then blocked with 10% normal goat serum (Sigma) in staining buffer. Sections prepared as above were first incubated for 18 h at 25°C with 1/500 dilution of polyclonal antibody or control rabbit serum in staining buffer. In the second step, the sections were incubated for 1 h at room temperature with biotin labeled goat anti-rabbit IgG (2 µg/ml) (Vector Laboratories) in staining

buffer and then for 30 min at room temperature with the Vectastain Elite ABC Kit (Vector Laboratories) in staining buffer. The sections were then rinsed in HBSS-HEPES without saponin. Immunoenzyme tissue staining was revealed with 3, 3' - diaminobenzidine tetrahydrochloride (DAB) substrate (0.5 mg/ml) (Sigma) in 0.05M Tris, pH 7.4, containing 0.0075% hydrogen peroxide. The substrate reaction was stopped by rinsing the sections in distilled water. The sections were then counterstained with Harris' hematoxylin (Shandon Lipshaw, Pittsburgh, PA).

The expression of TECK mRNA in murine adult thymus was analyzed by in situ hybridization and revealed a discrete positive non-lymphoid population within the thymus medulla. The expression of TECK protein was analyzed by using a polyclonal anti-serum made in a rabbit immunised with a peptide that consisted in the last 12 amino-acid of the murine TECK protein sequence. This polyclonal antibody reacts with the murine TECK recombinant protein prepared at DNAX both in ELISA and western blot. The application of this anti-serum on mouse adult thymic sections confirmed the distribution pattern obtained by in situ hybridization: the cells producing TECK are medullary stromal cells. The precise cell type producing TECK within the mouse thymus was identified, using the same anti-serum on sorted thymic subsets, as being the thymic dendritic cells.

R. Production of recombinant mouse TECK in *Escherichia coli* and other chemokines

Mouse recombinant TECK was produced in *E. coli* as a N-terminal FLAG (DYKDDDDKL; SEQ ID NO: 19) fusion protein. Briefly, the fusion construct containing FLAG followed by the mTECK sequence minus the leader peptide (see Table 1) was obtained by PCR amplification of the TECK cDNA in order to flank the coding sequence with HindIII and EcoRI sites and subsequent ligation in the pFLAG.1 vector which contains the FLAG sequence and an OmpA signal sequence. Electro-competent UT 4400 *E. coli* were transformed with the pFLAG.1-mTECK plasmid. The cells were grown in 2 x LB plus

50 µg/ml Ampicillin, induced at an OD. of 2.3 with 400 µM IPTG and harvested. The cell pellet was resuspended in cold lysis buffer (20 mM Tris pH 8, 2 mM EDTA, 20% sucrose, 0.1 mg/ml lysozyme, 100 µl Benzonase), homogenized and allowed to sit for 30 min. Then the same amount of a 1:4 dilution of cold lysis buffer without lysozyme was added for 10 more min. The solution was spun and the supernatant was filtered through a 0.2 µm membrane and then diluted 1:1 in 50 mM Tris pH 7.5. The diluted osmotic extract was submitted to chromatography on a Q-sepharose column equilibrated with 50 mM Tris pH 7.5 and eluted with a linear salt gradient. The fractions containing the recombinant protein were pooled. The fractions were then loaded onto a S-sepharose column equilibrated with 20 mM acetate pH 4.0. The column was eluted with a linear salt gradient and then with a 1.5M NaCl wash that contained the protein. Finally, the eluate was loaded onto a reverse phase column. The column was eluted with a linear gradient of 20% to 80% acetonitrile + 0.1% TFA. The concentration of the mTECK protein was estimated by Comassie blue staining and densitometric scanning of a 10% Nu-PAGE gel with lysozyme as a standard. The purity was estimated at 100% by sequencing of the N-terminus of the recombinant protein. Recombinant murine MIP-1α (R&D Systems, Minneapolis, MN) and lymphotactin (Hedrick, et al. (1997) J. Immunol. 158:1533-1540) were used as controls.

#### S. Assay for chemotaxis

The in vitro migration of cells isolated as described above in response to TECK or other factors was assessed in a modified Boyden micro chamber (Neuroprobe, Cabin John, MD) as described previously (Kelner, et al. (1994)). Briefly, factor dilutions in DMEM medium (Gibco) were loaded in the lower compartment in duplicate and 10<sup>5</sup> cells in a 50 µl volume of DMEM were loaded in the upper compartment. The two compartments were separated by a 5-µm or 8-µm pore size polycarbonate filter (Nucleopore, Pleasanton, CA). After incubation at 37° C for 80 min (or

120 min for lymphocytes), the filters were fixed in methanol and stained with Fields A and B. Cells migrated on the other side of the membrane were counted per five high-power fields (100 x) under microscope. The chemotactic index was calculated from the number of cells counted with the test sample divided by the number of cells counted with medium alone.

Northern blot analysis was performed of RNA from different organs hybridized with the mTECK cDNA probe with or without in vivo LPS stimulation. Hybridizing bands corresponded to the predicted  $\approx 1040$  bp size for mTECK mRNA. Significant induction occurred in spleen (with virtually no background), and in thymus and small intestine (both with higher background); no signal was detected in either condition for heart, lung, kidney, or liver.

mTECK mRNA expression was analysed in the mouse fetal thymus. RNA's from fetal thymic lobes were extracted at day 14, 15, 16 and 17 of gestation. Positive RT-PCR signals were detected in each of day 14, 15, 16, and 17 samples.

mTECK mRNA expression in thymic dendritic cells was evaluated. A population enriched in thymic dendritic cells was prepared from 15 pooled adult thymuses.  $>99\%$  pure dendritic cells were then sorted by flow cytometry based on their MHC Class II+ N-418+ phenotype. mTECK mRNA was then analyzed by RT-PCR and a MHC class II+ N-418- population sorted in the same experiment was used as a negative control. The N418+ sample gave a positive signal, while the N418- sample did not.

Expression analysis was performed with hTECK mRNA in different Human Tissues and Cell Types. Southern blots of human cDNA libraries digested with the appropriate restriction enzymes were hybridized with the hTECK cDNA probe. A major band hybridizing corresponding to the predicted length of hTECK mRNA ( $\approx 1040$  bp) was observed with sometimes some other bands that may represent incomplete cDNAs. Positive signals were detected in tonsil, fetal



spleen, and fetal small intestine. No signal was detected in activated (with PMA and ionomycin for 12 h) NK cells, activated (anti-CD40 antibody and IL-4 for 6 and 12 h) splenocytes,  $\gamma\delta$  T cells, activated (with anti-CD3 and PMA for 6, 12, and 24 h) PBMC, fetal testis, C+ (elutriated monocytes cultured with IFN- $\gamma$  and IL-10) monocytes, C-monocytes, 70% pure dendritic cells (CD1 $\alpha$ + dendritic cell population obtained by expansion of CD34+ bone marrow cells with GM-CSF and TNF- $\alpha$  and resting), and DC3 (similar dendritic cell population stimulated with PMA and ionomycin for 1 and 6 h), DC5 (dendritic cells obtained by culturing peripheral blood monocytes in the presence of IL-4 and GM-CSF), U937 (premonocytic cell line), and CD1 $\alpha$  cell lines. Ras KO mouse cDNA again confirmed that the mouse and human genes crosshybridize.

Four independent lines of transgenic mice expressing TECK in the brain have been made. All animals had neurologic disorders. In addition, several of them suffered severe infections. The consequences of TECK could be a direct one on brain cells which nature remains to be identify. Alternatively, since TECK has been shown in vitro to have effects on macrophages and dendritic cells which are critical effectors of immune responses, the overproduction of TECK could lead to distant effects on these cells at sites of infection. These results suggest that the blockade of TECK production in vivo may help to resolve particular pathological processes, in particular infections. The localization suggests a physiological role in immunological responses involving the thymus, or in colon/small intestine or gastrointestinal inflammation, e.g., Crohn's disease or inflammatory bowel disease.

## VII. Specific Characterization of the M/DC CR (CRAM)

Abbreviations: BAC, bacterial artificial chromosome; bp, base pair; CKR, chemokine receptor; EST, expressed sequence tag; GPR, G-protein-linked receptor; PBMC, peripheral blood mononuclear cells; STS, sequence tagged site.

We describe a novel human gene with high homology to CC- or  $\beta$ -chemokine receptors (CKRs). This putative CKR, CRAM, is most similar to human CCR1, with 46% amino acid identity and 65% similarity. CRAM is encoded by at least two alternatively spliced 1.5 and 1.8 kb mRNAs which specify at least two proteins differing by 12 amino acids at the N-terminus (CRAM-A and CRAM-B). CRAM mRNA was detected mainly in lymphoid tissues and expressed in activated monocytes, but not in B- or T-lymphocytes. CRAM mRNA expression was increased upon stimulation with IFN $\gamma$  and LPS but was not detectably inhibited by interleukin-10. CRAM was localized to the  $\beta$ -CKR cluster at chromosome 3p21 and physically linked to the CCR2 and CCR5 genes. In view of its similarity and genomic linkage to  $\beta$ -CKRs and restricted expression pattern, CRAM may play an important role in immune function. The existence of CRAM with alternative N-termini suggests a mechanism for altering ligand specificity and possibly signalling capacity of a single CKR.

Chemokines play critical roles in the chemoattraction and activation of leukocytes (Premack and Schall (1996) Nat Med 2:1174; Murphy (1996) Cytokine Growth Factor Rev 7:47; and Furie and Randolph (1995) Am J Pathol 146:1287), and have been divided into four families, based on the spacing of the first two of (usually) four conserved cysteine residues. The  $\alpha$  chemokines, with a C-X-C motif, include IL-8, MIP-2 $\alpha$ , GRO $\beta$ , and ENA-78. The  $\beta$  chemokines (C-C motif), include MIP-1 $\alpha$ , MCP-1, TARC, and RANTES. Recently, two new chemokine families have been defined by lymphotactin ( $\gamma$ ) and CX<sub>3</sub>Ckine ( $\delta$ ). Lymphotactin has only a

single cysteine residue at the corresponding location for the C-C or C-X-C motif. Kelner and Zlotnik. (1995) J Leukoc Biol 57:778; Kennedy, et al. (1995) J Immunol 155:203. CX<sub>3</sub>Ckine contains two cysteines separated by three  
5 intervening amino acids, and is tethered to the cell membrane via a long carboxy-terminal tail of mucin-like repeats. Bazan, et al. (1997) Nature 385:640.

Receptors for chemokines (CKRs) are G-protein coupled receptors (GPRs) with seven transmembrane domains. Novel  
10 CKRs have been identified by expression cloning of receptors binding a particular chemokine ligand (Holmes, et al. (1991) Science 253:1280) or mediating HIV fusion (Feng, et al. (1996) Science 272:872), by PCR using degenerate  
15 primers specific for conserved regions (Meyer, et al. (1996) J Biol Chem 271:14445; Ponath, et al. (1996) J Exp Med 183:2437; Daugherty, et al. (1996) J Exp Med 183:2349; Kurihara and Bravo (1996) J Biol Chem 271:11603; Power, et al. (1995) J Biol Chem 270:19495; Napolitano, et al. (1996) J Immunol 157:2759; and Raport, et al. (1996) J Leukoc Biol  
20 59:18), and by random sequencing efforts followed by sequence analysis. While nearly 30 CKR-like genes have been cloned from mammals and mammalian viruses, only 17 have been shown to bind identified chemokines. Thus, a substantial number of CKR-like molecules remain "orphan  
25 receptors." Most CKRs with experimentally identified ligands bind to more than one ligand. IL-8 receptor B (CXCR2) binds to the  $\alpha$  chemokines IL-8, NAP-2, and MGSA (Suzuki, et al. (1994) J Biol Chem 269:18263), whereas human CCR5 binds the  $\beta$  chemokines RANTES, MIP-1 $\alpha$ , and MIP-  
30 1 $\beta$  (Raport, et al. (1996) J Biol Chem 271:17161; and Alkhatib, et al. (1996) Science 272:1955).

We have used cDNA library subtraction to isolate genes which are induced by monocyte activation. We thereby  
isolated a cDNA clone from a subtracted library enriched  
35 for monocyte activation-specific cDNAs that shows considerable homology to CC- or  $\beta$ -CKRs and maps within the  $\beta$ -CKR cluster on human chromosome 3p21. Expression of this

gene was detected in several lymphoid tissues and in activated monocytes (but not lymphocytes). We provisionally designate this gene CRAM, for chemokine receptor of activated monocytes. CRAM is expressed as at least two alternatively spliced mRNAs encoding CKRs with different N-terminal amino acid sequences, suggesting a possible novel mechanism for regulation of CKR ligand specificity.

#### A. Cell cultures and cDNA library construction

Human PBMC were purified by density gradient centrifugation on Ficoll (Pharmacia Biotech Inc., Piscataway, NJ) using standard procedures. Monocytes were enriched from PBMC by adherence to tissue culture flasks and cultured in DMEM + 10% FCS. Monocytes were activated by culture with 100 ng/ml IFN $\gamma$  (R & D Systems Inc., Minneapolis, MN) and 1  $\mu$ g/ml LPS (Life Technologies, Grand Island, NY) for 1-15 hr. Total RNA was prepared by guanidinium isothiocyanate lysis followed by poly(A)+ RNA selection using the OLIGOTEX kit (QIAGEN Inc., Chatsworth, CA). cDNA libraries containing  $>2 \times 10^6$  independent clones were constructed using the SuperScript cDNA Kit (Life Technologies).

#### B. cDNA library subtractions

Subtracted cDNA libraries (activated monocytes minus resting PBMC) were constructed. See, e.g., Hara, et al. (1994) Blood 84:189; and Kennedy, et al. (1996) J Interferon Cytokine Res 16:611. The major cDNA species present in the subtracted library were then added (1  $\mu$ g each) to the resting PBMC cDNA library (150  $\mu$ g); this mixture was used as the driver cDNA for a second round of subtraction using 5  $\mu$ g of the activated monocyte cDNA library to enrich for induction-specific cDNAs which were less abundantly expressed.

#### C. DNA sequencing and bioinformatics

The nucleotide sequence of CRAM was determined using an ABI 377 automated sequencer and standard techniques. DNA sequence analyses were performed using Sequencher 3.0

(Gene Codes Corporation, Ann Arbor, MI) and MacVector 6.0 (Oxford Molecular Group). Comparisons to GenBank databases were performed using the BLAST program on web-based servers: (<http://www.ncbi.nlm.nih.gov/BLAST/> and

- 5 <http://www.genome.ad.jp/SIT/BLAST.html>). Sequence alignments and phylogenetic analyses utilized ClustalW 1.6 (Higgins, et al. (1996) Methods in Enzymology 266:383) and TreeViewPPC 1.2 (Page (1996) Computer Applications in the Biosciences 12:357).

10 D. Analysis of CRAM mRNA expression

- Multiple-tissue Northern blots were purchased from Clontech (Palo Alto, CA). Poly(A)+ RNA from human monocytes was used for RNA blot analysis. cDNA libraries from human cells (5 µg) in the pSPORT vector (Life
- 15 Technologies) were digested with SalI and NotI to release cDNA inserts, electrophoresed on 1% agarose gels, and subjected to Southern blot transfer/hybridization. Hybridizations with <sup>32</sup>P-labeled CRAM DNA fragments encoding the C-terminal 144 amino acids of the predicted ORF were
- 20 done at 65° C in ExpressHyb (Clontech, Palo Alto, CA) for 2 hr, followed by two stringent washes at 50° C in 0.1X SSC, 0.1% SDS for 45 min. Hybridization was detected using a STORM 860 phosphorimager (Molecular Dynamics, Sunnyvale, CA). Reverse transcriptase PCR (RT-PCR) was performed with
- 25 Superscript II reverse transcriptase (Life Technologies) and Taq DNA polymerase (Boehringer-Mannheim, Indianapolis, IN). PCR was for 35 cycles of 95° C/45 sec, 62° C/30 sec, 72° C/60 sec. Primers specific for exon 1 (5'-AGACGCTTCAGAGATCCTCTGGAGGCC) or exon 2 (5'-GAAGCTGCTTCGGGGGTGAGCAAAC) were used in conjunction
- 30 with an exon 3-specific primer (5'-CAAACACAGCAGAGCAGAGTGATGGCACC) for amplification.

E. Chromosomal localization

- PCR was performed on genomic DNA from the 83 cell
- 35 lines of the Stanford Human Genome Center G3 radiation hybrid panel (Research Genetics, Huntsville, AL) using CRAM primers: (5'-GTGTCCTGGCATGGGTAACAGCC) and

(5'-CGGTGGAATGGTCAGGTTCTTCCC) as previously described for the GeneBridge 4 radiation hybrid panel (Samson, et al. (1996) Genomics 36:522). Data correlating the presence or absence of PCR product to each cell line were entered into the RHserver (Stanford Human Genome Center; <http://shgc.stanford.edu/RH/>). Co-localized STSs were identified on the human physical map using the Entrez server (National Center for Biotechnology Information; <http://www3.ncbi.nlm.nih.gov/Entrez/>).

#### 10 F. cDNA cloning of CRAM

We employed subtractive hybridization to identify genes induced in monocytes upon activation by IFN $\gamma$  and LPS. An activated monocyte cDNA library was first subtracted against a resting PBMC cDNA library. Seven prominent  
15 induced cDNAs thus identified were mixed with the resting PBMC library, which was then used as "driver" in another subtraction to generate a new library containing less abundantly expressed, induction-specific cDNAs. More than  
20 100 clones were isolated from this second-round subtracted library, representing 55 unique cDNAs, 25 of which did not correspond to known cDNAs from the non-redundant section of GenBank. One of these clones contained a 1.5 kb insert encoding a large open reading frame with strong homology to all five known human  $\beta$ -CKRs. We designated this cDNA CRAM  
25 (chemokine receptor of activated monocytes; or M/DC CR).

#### G. Sequence analysis of CKRs

A phylogenetic analysis of CKRs and related gene sequences revealed two major clades or phylogenetic groups, with several receptors remaining unclustered outside these  
30 two groups. Interestingly, the two groups correlated with known ligand specificities: the  $\alpha$ -CKR IL-8RA, IL-8RB, and fusin cluster in a single clade, while  $\beta$ -CKR CCR1 through CCR5 all cluster in a second clade. Of the seven receptors that do not fall into either group, one (DARC) is a  
35 promiscuous CKR that binds several  $\alpha$ - and  $\beta$ -chemokines (Neote, et al. (1993) J Biol Chem 268:12247).

The 1536 bp CRAM cDNA encodes an ORF with a predicted size of about 356 amino acids. Phylogenetic analysis showed that CRAM was most closely related to  $\beta$ -CKRs, exhibiting strongest homology to CCR1 (46% identity and 65% similarity), and the least to CCR4, with only 33% identity and 48% similarity. Three other human orphan receptors V28, TER1, and GPR5 also group with  $\beta$ -CKRs, and like CRAM, may be receptors for known or yet to be identified  $\beta$  chemokines.

10       The two most highly conserved regions among CCR1 through CCR5 are in transmembrane region 2 (YLLNLAISDLLF; "TM2") and immediately after transmembrane region 3 (IDRYLAIVHAVF; "DRY-box"). These two 12-amino acid segments are invariant among CCR1 through CCR4; CCR5 shares 15 22 of these 24 residues. These regions are sometimes conserved among other mammalian GPR and have been used for degenerate primer PCR to clone new CKRs. CRAM is divergent in these regions (9 out of 12 amino acids in TM2; 4 out of 12 in the DRY-box), which may explain why such approaches 20 have failed to identify CRAM. The DRY-box is in one of the three intracellular loops, and is thought to play a role in binding to heterotrimeric G proteins (Damaj, et al. (1996) FASEB J 10:1426). Because of the divergence of CRAM from the other  $\beta$ -CKRs in these regions, it may interact with a 25 different subset of G protein subunits, possibly transducing a signal different from that induced via other  $\beta$ -CKRs.

While human CKR genes have been localized to several different chromosomes, the  $\beta$ -CKR genes CCR1, CCR2, CCR3, 30 and CCR5 all cluster in a 350 kb region at chromosome 3p21.3 (Samson, et al. (1996) Genomics 36:522). CCR4 and the orphan receptors TER1 and GPR5 are also located in this 3p21 region Napolitano, et al. (1996) J Immunol 157:2759; Samson, et al. (1996) Genomics 36:522; Heiber, et al. 35 (1995) DNA Cell Biol 14:25) We determined the chromosomal location of CRAM. The Stanford G3 panel of radiation hybrids was used as templates for PCR reactions with CRAM-

specific primers. Among the 83 different hybrids, 11 contained the CRAM gene as assessed by PCR. CRAM co-localized with STS D3S3888, which is located at chromosome 3p21.3. Confirmation of this result was obtained from the recently completed sequence of the 143 kb BAC clone 110p12 from the 3p21 region (GenBank accession U95626); this BAC contains the loci CCR2, CCR5, and CRAM.

A related but different CRAM cDNA was also isolated from an activated monocyte library by random sequencing. Comparing the two forms of CRAM to the genomic sequence revealed the existence of two short exons (corresponding to 95750-96064 bp and 96186-96256 bp on BAC 110p12), followed by a large third exon (96630-98093 bp) that contains almost the entire ORF for CRAM. These two CRAM cDNAs consist of either exon 2 and exon 3 (1536 bp), or exon 1 and exon 3 (1780 bp). Exon 2 contributes 12 amino acids in frame with exon 3 to form the entire 356 residue polypeptide (CRAM-A). As exon 1 has no methionine in frame with the ORF in exon 3, the translated protein from this splice variant would start with Met-13, resulting in an N-terminally truncated protein of 344 amino acids (CRAM-B).

#### H. Expression of CRAM mRNA

RNA blot analysis showed expression largely restricted to lymphoid tissues. Prominent expression of CRAM mRNA was observed in spleen, lymph node, thymus, bone marrow, and fetal liver. Very little expression was detected in brain, liver, muscle, kidney, pancreas, or PBL, with moderate signals in heart, placenta, lung, and appendix. This pattern of expression was similar to that of the CKR-like gene *TER1* (Napolitano, et al. (1996) J Immunol 157:2759), but quite different from the related orphan receptor genes *V28* and *blr1* (Forster, et al. (1996) Cell 87:1037; and Raport, et al. (1995) Gene 163:295).

Data from various hematopoietic cell types showed no evidence for CRAM expression in resting or activated lymphocytes, or in splenocytes. CRAM mRNA was also not detected in resting monocytic cell lines, but was strongly



expressed in primary monocytes and THP-1 cells upon activation with IFN $\gamma$  and LPS. Both CRAM-A and CRAM-B mRNAs were induced, as detected by RT-PCR using exon 1- and exon 2-specific primers. In contrast to several other monocyte activation-induced genes, such as monokines (TNF $\alpha$ , IL-1, IL-6) and some cell-surface antigens (Ho and Moore. (1994) Therapeutic Immunology 1:173). CRAM mRNA expression was not detectably inhibited by IL-10. Thus, CRAM expression in monocytes may be regulated via a different mechanism compared to that of several other activation-induced genes.

While most CKR genes lack introns, the genes for human CCR2 and mouse CXCR4 (fusin) both contain at least two exons and both have two alternatively spliced forms. CCR2A and CCR2B differ in the C-terminus (Charo, et al. (1994) Proc. Natl. Acad. Sci. U.S.A. 91:2752), whereas CXCR4 has two forms that differ by two amino acids at the N-terminus (Heesen, et al. (1997) J. Immunol. 158:3561). The two forms of CCR2 have identical ligand specificities, but differ with respect to which G $\alpha$  subunits they can couple (Kuang, et al. (1996) J. Biol. Chem. 271:3975); the two forms of CXCR4 can both serve as functional CKRs for SDF-1 $\alpha$  (Heesen, et al. (1997) J. Immunol. 158:3561), although their ligand specificities and interactions with HIV have not been fully characterized. The N-terminal sequence of CKRs, along with portions of the extracellular loops, is known to play a key role in ligand binding and possibly receptor activation (Ahuja, et al. (1996) J. Biol. Chem. 271:225; Lu, et al. (1995) J. Biol. Chem. 270:26239; Horuk (1994) Immunol. Today 15:169; Wells, et al. (1996) J Leukoc Biol 59:53; and Hebert, et al. (1993) J. Biol. Chem. 268:18549). This region of CKR is also important for HIV fusion (Rucker, et al. (1996) Cell 87:437.), which is antagonized by chemokine ligands (Paxton, et al. (1996) Nature Med. 2:412; and Cocchi, et al. (1995) Science 270:1811). Thus it is possible that CRAM-A and CRAM-B may exhibit different but likely overlapping ligand specificities. Regulated expression of alternative forms

of a single CKR, combined with possible modulation of specificity of ligand-receptor interaction by chemokine-proteoglycan interaction (Graham, et al. (1996) The EMBO J. 15:6506; and Witt and Lander (1994) Curr. Biol. 4:394),  
5 might control the spectrum of chemokines to which a particular cell could respond. In addition, these observations may provide one possible explanation of non-reciprocal desensitization phenomena observed with, for example, the chemokines RANTES, MIP-1 $\alpha$ , and MCAF (Wang, et  
10 al. (1993) J Exp Med 177:699).

The similarity of CRAM to the other  $\beta$ -CKRs, its chromosomal localization in the  $\beta$ -CKR gene cluster, and induction of its expression in monocytes upon activation all argue that CRAM may play an important role in  
15 regulation of immune function.

#### VIII. Screening for receptor/ligand

Labeled reagent is useful for screening of an  
20 expression library made from a cell line which expresses a chemokine or receptor, as appropriate. Standard staining techniques are used to detect or sort intracellular or surface expressed ligand, or surface expressing transformed cells are screened by panning. Screening of intracellular  
25 expression is performed by various staining or immunofluorescence procedures. See also, e.g., McMahan, et al. (1991) EMBO J. 10:2821-2832.

For example, on day 0, precoat 2-chamber permanox slides with 1 ml per chamber of fibronectin, 10 ng/ml in  
30 PBS, for 30 min at room temperature. Rinse once with PBS. Then plate COS cells at  $2-3 \times 10^5$  cells per chamber in 1.5 ml of growth media. Incubate overnight at 37° C.

On day 1 for each sample, prepare 0.5 ml of a solution of 66  $\mu$ g/ml DEAE-dextran, 66  $\mu$ M chloroquine, and 4  $\mu$ g DNA  
35 in serum free DME. For each set, a positive control is prepared, e.g., of huIL-10-FLAG cDNA at 1 and 1/200 dilution, and a negative mock. Rinse cells with serum free

DME. Add the DNA solution and incubate 5 hr at 37° C. Remove the medium and add 0.5 ml 10% DMSO in DME for 2.5 min. Remove and wash once with DME. Add 1.5 ml growth medium and incubate overnight.

- 5 On day 2, change the medium. On days 3 or 4, the cells are fixed and stained. Rinse the cells twice with Hank's Buffered Saline Solution (HBSS) and fix in 4% paraformaldehyde (PFA)/glucose for 5 min. Wash 3X with HBSS. The slides may be stored at -80° C after all liquid
- 10 is removed. For each chamber, 0.5 ml incubations are performed as follows. Add HBSS/saponin(0.1%) with 32 µl/ml of 1M NaN<sub>3</sub> for 20 min. Cells are then washed with HBSS/saponin 1X. Add antibody complex to cells and incubate for 30 min. Wash cells twice with HBSS/saponin.
- 15 Add second antibody, e.g., Vector anti-mouse antibody, at 1/200 dilution, and incubate for 30 min. Prepare ELISA solution, e.g., Vector Elite ABC horseradish peroxidase solution, and preincubate for 30 min. Use, e.g., 1 drop of solution A (avidin) and 1 drop solution B (biotin) per 2.5
- 20 ml HBSS/saponin. Wash cells twice with HBSS/saponin. Add ABC HRP solution and incubate for 30 min. Wash cells twice with HBSS, second wash for 2 min, which closes cells. Then add Vector diaminobenzoic acid (DAB) for 5 to 10 min. Use 2 drops of buffer plus 4 drops DAB plus 2 drops of H<sub>2</sub>O<sub>2</sub> per
- 25 5 ml of glass distilled water. Carefully remove chamber and rinse slide in water. Air dry for a few minutes, then add 1 drop of Crystal Mount and a cover slip. Bake for 5 min at 85-90° C.

- Alternatively, the binding compositions are used to
- 30 affinity purify or sort out cells expressing the ligand or receptor. See, e.g., Sambrook, et al. or Ausubel et al.

- All references cited herein are incorporated herein by reference to the same extent as if each individual publication or patent application was specifically and
- 35 individually indicated to be incorporated by reference.

Many modification an variations of this invention can be made without departing from its spirit and scope, as

will be apparent to those skilled in the art. The specific embodiments described herein are offered by way of example only, and the invention is to be limited only by the terms of the appended claims, along with the full scope of the  
5 equivalents to which such claims are entitled.

## SEQUENCE LISTING

- 5 SEQ ID NO: 1 is mouse TECK nucleotide sequence.  
SEQ ID NO: 2 is mouse TECK amino acid sequence.  
SEQ ID NO: 3 is human TECK nucleotide sequence.  
SEQ ID NO: 4 is human TECK amino acid sequence.  
SEQ ID NO: 5 is human MIP-3 $\alpha$  nucleotide sequence.  
SEQ ID NO: 6 is human MIP-3 $\alpha$  amino acid sequence.  
10 SEQ ID NO: 7 is human MIP-3 $\beta$  nucleotide sequence.  
SEQ ID NO: 8 is human MIP-3 $\beta$  amino acid sequence.  
SEQ ID NO: 9 is human DC CR nucleotide sequence.  
SEQ ID NO: 10 is human DC CR amino acid sequence.  
SEQ ID NO: 11 is human M/DC CR nucleotide sequence.  
15 SEQ ID NO: 12 is human M/DC CR amino acid sequence.  
SEQ ID NO: 13 is human CCKR1 amino acid sequence.  
SEQ ID NO: 14 is human CCKR2 amino acid sequence.  
SEQ ID NO: 15 is human CCKR3 amino acid sequence.  
SEQ ID NO: 16 is human CCKR4 amino acid sequence.  
20 SEQ ID NO: 17 is HPRT sense primer.  
SEQ ID NO: 18 is HPRT antisense primer.  
SEQ ID NO: 19 is FLAG epitope tag sequence.
- 25 (1) GENERAL INFORMATION:
- (i) APPLICANT: Wang, Wei  
Gish, Kurt C.  
Schall, Thomas J.  
30 Vicari, Alain P.  
Zlotnik, Albert
- (ii) TITLE OF INVENTION: MAMMALIAN CHEMOKINE REAGENTS
- 35 (iii) NUMBER OF SEQUENCES: 19
- (iv) CORRESPONDENCE ADDRESS:  
(A) ADDRESSEE: DNAX Research Institute  
(B) STREET: 901 California Avenue  
40 (C) CITY: Palo Alto  
(D) STATE: California  
(E) COUNTRY: USA  
(F) ZIP: 94304-1104
- 45 (v) COMPUTER READABLE FORM:  
(A) MEDIUM TYPE: Floppy disk  
(B) COMPUTER: IBM PC compatible  
(C) OPERATING SYSTEM: PC-DOS/MS-DOS  
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30
- 50 (vi) CURRENT APPLICATION DATA:  
(A) APPLICATION NUMBER:  
(B) FILING DATE:  
(C) CLASSIFICATION:
- 55 (vii) PRIOR APPLICATION DATA:  
(A) APPLICATION NUMBER: US 08/675,814  
(B) FILING DATE: 05-JUL-1996
- 60 provisional filings DX0589P, DX0589P1; DX0589P2

various

## (viii) ATTORNEY/AGENT INFORMATION:

- (A) NAME: Ching, Edwin P.  
(B) REGISTRATION NUMBER: 34,090  
(C) REFERENCE/DOCKET NUMBER: DX0589Q1

## (ix) TELECOMMUNICATION INFORMATION:

- (A) TELEPHONE: 415-852-9196  
(B) TELEFAX: 415-496-1200

## (2) INFORMATION FOR SEQ ID NO:1:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1034 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: cDNA

## (ix) FEATURE:

- (A) NAME/KEY: CDS  
(B) LOCATION: 94..525

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

30 AGGCTACAAG CAGGCACCAG CTCTCAGGAC CAGAAAGGCA TTGGTGGCCC CCTTAAACCT 60  
TCAGGTATCT GGAGAGGAGA TCTAACCTTC ACT ATG AAA CTG TGG CTT TTT GCC 114  
Met Lys Leu Trp Leu Phe Ala  
1 5  
35 TGC CTG GTT GCC TGT TTT GTT GGG GCC TGG ATG CCG GTT GTC CAT GCC 162  
Cys Leu Val Ala Cys Phe Val Gly Ala Trp Met Pro Val Val His Ala  
10 15 20  
40 CAA GGT GCC TTT GAA GAC TGC TGC CTG GGT TAC CAG CAC AGG ATC AAA 210  
Gln Gly Ala Phe Glu Asp Cys Cys Leu Gly Tyr Gln His Arg Ile Lys  
25 30 35  
45 TGG AAT GTT CTC CGG CAT GCT AGG AAT TAT CAC CAG CAG GAA GTG AGT 258  
Trp Asn Val Leu Arg His Ala Arg Asn Tyr His Gln Gln Glu Val Ser  
40 45 50 55  
50 GGA AGC TGC AAC CTA CGT GCT GTG AGA TTC TAC TTC CGC CAG AAA GTA 306  
Gly Ser Cys Asn Leu Arg Ala Val Arg Phe Tyr Phe Arg Gln Lys Val  
60 65 70  
55 GTG TGT GGG AAT CCA GAG GAC ATG AAT GTG AAG AGG GCG ATA AGA ATC 354  
Val Cys Gly Asn Pro Glu Asp Met Asn Val Lys Arg Ala Ile Arg Ile  
75 80 85  
60 TTG ACA GCT AGG AAA AGG CTA GTC CAC TGG AAG AGC GCC TCA GAC TCT 402  
Leu Thr Ala Arg Lys Arg Leu Val His Trp Lys Ser Ala Ser Asp Ser  
90 95 100  
60 CAG ACT GAA AGG AAG AAG TCA AAC CAT ATG AAG TCC AAG GTG GAG AAC 450

Gln Thr Glu Arg Lys Lys Ser Asn His Met Lys Ser Lys Val Glu Asn  
 105 110 115

5 CCC AAC AGT ACA AGC GTG AGG AGT GCC ACC CTA GGT CAT CCC AGG ATG 498  
 Pro Asn Ser Thr Ser Val Arg Ser Ala Thr Leu Gly His Pro Arg Met  
 120 125 130 135

10 GTG ATG ATG CCC AGA AAG ACC AAC AAT TAAGTTAATT ACTCAGAGTA 545  
 Val Met Met Pro Arg Lys Thr Asn Asn  
 140

15 AGCACCAGCT GGAGGATGGG CGGAGTCTGC TGAAGTGCTG TCTTCTAGGC ATGCCAGTGC 605  
 CAATGAACTC ACTGAAGCTA CAGTTTCCTG TACAAGACCA GACCCACCAA CGTCTCAGCA 665  
 TGTACGAGGA AGGAACTACT GCGCTAAAGG CCCTCCCACT CACCAAGGAG CTATTGGCTA 725  
 TTGATGATTG CTGAGGGAAG GGAGTAATTT TTTTCTCTT TCTGAAGTGT GACTTGAGTA 785

20 AATTGCCCCAT AGTTCAGTAT ATAATCCCCA ACCTGTGCTC AGGCAAGCAA CCCTAATTAA 845  
 ATGCAATAGC CACATACAAA AGAAGAGGAT ATGAATAGTT TGGTAGGAGG GGCTTGTTAG 905  
 GAAGAAGACA TTAACAGGAG AGAGAGGAGC GAGAGGATAG TGAGTGTGTG AGAGTGCCTG 965

25 CACGTGTGAA ATGGTCAAAG AATTAAAAA TAAAACTTA AAAAGCTATT AAAAAGTAAA 1025  
 AAAAATAAA 1034

30 (2) INFORMATION FOR SEQ ID NO:2:

35 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 144 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

40 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Lys Leu Trp Leu Phe Ala Cys Leu Val Ala Cys Phe Val Gly Ala  
 1 5 10 15

45 Trp Met Pro Val Val His Ala Gln Gly Ala Phe Glu Asp Cys Cys Leu  
 20 25 30

Gly Tyr Gln His Arg Ile Lys Trp Asn Val Leu Arg His Ala Arg Asn  
 35 40 45

50 Tyr His Gln Gln Glu Val Ser Gly Ser Cys Asn Leu Arg Ala Val Arg  
 50 55 60

55 Phe Tyr Phe Arg Gln Lys Val Val Cys Gly Asn Pro Glu Asp Met Asn  
 65 70 75 80

Val Lys Arg Ala Ile Arg Ile Leu Thr Ala Arg Lys Arg Leu Val His  
 85 90 95

60 Trp Lys Ser Ala Ser Asp Ser Gln Thr Glu Arg Lys Lys Ser Asn His

100 105 110

Met Lys Ser Lys Val Glu Asn Pro Asn Ser Thr Ser Val Arg Ser Ala  
 115 120 125

5 Thr Leu Gly His Pro Arg Met Val Met Met Pro Arg Lys Thr Asn Asn  
 130 135 140

## 10 (2) INFORMATION FOR SEQ ID NO:3:

## (i) SEQUENCE CHARACTERISTICS:

- 15 (A) LENGTH: 1012 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: cDNA

## 20 (ix) FEATURE:

- (A) NAME/KEY: CDS  
 (B) LOCATION: 117..566

## 25 (ix) FEATURE:

- (A) NAME/KEY: mat\_peptide  
 (B) LOCATION: 186..566

## 30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

TCGACCCACG CGTCCGCTTG GCCTACAGCC CGGCGGGCAT CAGCTCCCTT GACCCAGTGG 60

ATATCGGTGG CCCC GTTATT CGTCCAGGTG CCCAGGGAGG AGGACCCGCC TGCAGC 116

35 ATG AAC CTG TGG CTC CTG GCC TGC CTG GTG GCC GGC TTC CTG GGA GCC 164  
 Met Asn Leu Trp Leu Leu Ala Cys Leu Val Ala Gly Phe Leu Gly Ala  
 -23 -20 -15 -10

40 TGG GCC CCC GCT GTC CAC ACC CAA GGT GTC TTT GAG GAC TGC TGC CTG 212  
 Trp Ala Pro Ala Val His Thr Gln Gly Val Phe Glu Asp Cys Cys Leu  
 -5 1 5

GCC TAC CAC TAC CCC ATT GGG TGG GCT GTG CTC CGG CGC GCC TGG ACT 260  
 45 Ala Tyr His Tyr Pro Ile Gly Trp Ala Val Leu Arg Arg Ala Trp Thr  
 10 15 20 25

TAC CGG ATC CAG GAG GTG AGC GGG AGC TGC AAT CTG CCT GCT GCG ATA 308  
 50 Tyr Arg Ile Gln Glu Val Ser Gly Ser Cys Asn Leu Pro Ala Ala Ile  
 30 35 40

TTC TAC CTC CCC AAG AGA CAC AGG AAG GTG TGT GGG AAC CCC AAA AGC 356  
 Phe Tyr Leu Pro Lys Arg His Arg Lys Val Cys Gly Asn Pro Lys Ser  
 45 50 55

55 AGG GAG GTG CAG AGA GCC ATG AAG CTC CTG GAT GCT CGA AAT AAG GTT 404  
 Arg Glu Val Gln Arg Ala Met Lys Leu Leu Asp Ala Arg Asn Lys Val  
 60 65 70

60 TTT GCA AAG CTC CAC CAC AAC ATG CAG ACC TTC CAA GCA GGC CCT CAT 452



Phe Ala Lys Leu His His Asn Met Gln Thr Phe Gln Ala Gly Pro His  
 75 80 85

5 GCT GTA AAG AAG TTG AGT TCT GGA AAC TCC AAG TTA TCA TCA TCC AAG 500  
 Ala Val Lys Lys Leu Ser Ser Gly Asn Ser Lys Leu Ser Ser Ser Lys  
 90 95 100 105

10 TTT AGC AAT CCC ATC AGC AGC AGC AAG AGG AAT GTC TCC CTC CTG ATA 548  
 Phe Ser Asn Pro Ile Ser Ser Ser Lys Arg Asn Val Ser Leu Leu Ile  
 110 115 120

TCA GCT AAT TCA GGA CTG TGAGCCGGCT CATTTCTGGG CTCCATCGGC 596  
 Ser Ala Asn Ser Gly Leu  
 125

15 ACAGGAGGGG CCGGATCTTT CTCCGATAAA ACCGTCGCCC TACAGACCCA GCTGTCCCCA 656  
 CGCCTCTGTC TTTTGGGTCA AGTCTTAATC CCTGCACCTG AGTTGGTCCT CCCTCTGCAC 716

20 CCCCACCACC TCCTGCCCCG CTGGCAACTG GAAAGAAGGA GTTGGCCTGA TTTTAACCTT 776  
 TTGCCGCTCC GGGGAACAGC ACAATCCTGG GCAGCCAGTG GCTCTTGTAG AGAAAACTTA 836  
 GGATACCTCT CTCACCTTCT GTTTCTTGCC GTCCACCCCG GGCCATGCCA GTGTGTCCTC 896

25 TGGGTCCCCT CCAAAAATCT GGTCATTCAA GGATCCCCTC CCAAGGCTAT GCTTTTCTAT 956  
 AACTTTTAAA TAAACCTTGG GGGGTGAATG GAATAAAAAA AAAAAAAAAA AAAAAA 1012

30 (2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 150 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

35 (ii) MOLECULE TYPE: protein

40 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met Asn Leu Trp Leu Leu Ala Cys Leu Val Ala Gly Phe Leu Gly Ala  
 -23 -20 -15 -10

45 Trp Ala Pro Ala Val His Thr Gln Gly Val Phe Glu Asp Cys Cys Leu  
 -5 1 5

Ala Tyr His Tyr Pro Ile Gly Trp Ala Val Leu Arg Arg Ala Trp Thr  
 10 15 20 25

50 Tyr Arg Ile Gln Glu Val Ser Gly Ser Cys Asn Leu Pro Ala Ala Ile  
 30 35 40

55 Phe Tyr Leu Pro Lys Arg His Arg Lys Val Cys Gly Asn Pro Lys Ser  
 45 50 55

Arg Glu Val Gln Arg Ala Met Lys Leu Leu Asp Ala Arg Asn Lys Val  
 60 65 70

60 Phe Ala Lys Leu His His Asn Met Gln Thr Phe Gln Ala Gly Pro His

75 80 85

Ala Val Lys Lys Leu Ser Ser Gly Asn Ser Lys Leu Ser Ser Ser Lys  
 90 95 100 105

5 Phe Ser Asn Pro Ile Ser Ser Ser Lys Arg Asn Val Ser Leu Leu Ile  
 110 115 120

10 Ser Ala Asn Ser Gly Leu  
 125

(2) INFORMATION FOR SEQ ID NO:5:

15 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 801 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

20 (ii) MOLECULE TYPE: cDNA

(ix) FEATURE:  
 (A) NAME/KEY: CDS  
 25 (B) LOCATION: 1..288

(ix) FEATURE:  
 (A) NAME/KEY: mat\_peptide  
 30 (B) LOCATION: 79..288

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

35 ATG TGC TGT ACC AAG AGT TTG CTC CTG GCT GCT TTG ATG TCA GTG CTG 48  
 Met Cys Cys Thr Lys Ser Leu Leu Leu Ala Ala Leu Met Ser Val Leu  
 -26 -25 -20 -15

40 CTA CTC CAC CTC TGC GGC GAA TCA GAA GCA GCA AGC AAC TTT GAC TGC 96  
 Leu Leu His Leu Cys Gly Glu Ser Glu Ala Ala Ser Asn Phe Asp Cys  
 -10 -5 1 5

45 TGT CTT GGA TAC ACA GAC CGT ATT CTT CAT CCT AAA TTT ATT GTG GGC 144  
 Cys Leu Gly Tyr Thr Asp Arg Ile Leu His Pro Lys Phe Ile Val Gly  
 10 15 20

50 TTC ACA CGG CAG CTG GCC AAT GAA GGC TGT GAC ATC AAT GCT ATC ATC 192  
 Phe Thr Arg Gln Leu Ala Asn Glu Gly Cys Asp Ile Asn Ala Ile Ile  
 25 30 35

55 TTT CAC ACA AAG AAA AAG TTG TCT GTG TGC GCA AAT CCA AAA CAG ACT 240  
 Phe His Thr Lys Lys Lys Leu Ser Val Cys Ala Asn Pro Lys Gln Thr  
 40 45 50

60 TGG GTG AAA TAT ATT GTG CGT CTC CTC AGT AAA AAA GTC AAG AAC ATG 288  
 Trp Val Lys Tyr Ile Val Arg Leu Leu Ser Lys Lys Val Lys Asn Met  
 55 60 65 70

TAAAACTGT GGCTTTTCTG GAATGGAATT GGACATAGCC CAAGAACAGA AAGAACCTTG 348

60 CTGGGGTTGG AGGTTTCACT TGCACATCAT GGAGGGTTTA GTGCTTATCT AATTTGTGCC 408

5 TCACTGGACT TGTCCAATTA ATGAAGTTGA TTCATATTGC ATCATAGTTT GCTTTGTTTA 468  
AGCATCACAT TAAAGTTAAA CTGTATTTTA TGTTATTTAT AGCTGTAGGT TTTCTGTGTT 528  
TAGCTATTTA ATACTAATTT TCCATAAGCT ATTTTGTTT AGTGCAAAGT ATAAAATTAT 588  
ATTTGGGGGG GAATAAGATT ATATGGACTT TTTTGCAAGC AACAAGCTAT TTTTAAAAA 648  
10 AAACATTTTA ACATTCTTTT GTTTATATTG TTTTGTCTCC TAAATTGTTG TAATTGCATT 708  
ATAAAATAAG AAAAATATTA ATAAGACAAA TATTGAAAAT AAAGAAACAA AAAGTTAAAA 768  
AAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAA 801

(2) INFORMATION FOR SEQ ID NO:6:

20 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 96 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Met Cys Cys Thr Lys Ser Leu Leu Leu Ala Ala Leu Met Ser Val Leu  
-26 -25 -20 -15  
30 Leu Leu His Leu Cys Gly Glu Ser Glu Ala Ala Ser Asn Phe Asp Cys  
-10 -5 1 5  
Cys Leu Gly Tyr Thr Asp Arg Ile Leu His Pro Lys Phe Ile Val Gly  
35 10 15 20  
Phe Thr Arg Gln Leu Ala Asn Glu Gly Cys Asp Ile Asn Ala Ile Ile  
25 30 35  
40 Phe His Thr Lys Lys Lys Leu Ser Val Cys Ala Asn Pro Lys Gln Thr  
40 45 50  
Trp Val Lys Tyr Ile Val Arg Leu Leu Ser Lys Lys Val Lys Asn Met  
45 55 60 65 70

(2) INFORMATION FOR SEQ ID NO:7:

50 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 699 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

55 (ii) MOLECULE TYPE: cDNA

(ix) FEATURE:  
60 (A) NAME/KEY: CDS  
(B) LOCATION: 142..435

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

5 GGCACGAGCG GCACGAGCAT CACTCACACC TTGCATTTCA CCCCTGCATC CCAGTCGCCC 60  
 TGCAGCCTCA CACAGATCCT GCACACACCC AGACAGCTGG CGCTCACACA TTCACCGTTG 120  
 GCCTGCCTCT GTTCACCCTC C ATG GCC CTG CTA CTG GCC CTC AGC CTG CTG 171  
 10 Met Ala Leu Leu Leu Ala Leu Ser Leu Leu 10  
 1 GTT CTC TGG ACT TCC CCA GCC CCA ACT CTG AGT GGC ACC AAT GAT GCT 219  
 Val Leu Trp Thr Ser Pro Ala Pro Thr Leu Ser Gly Thr Asn Asp Ala 25  
 15 GAA GAC TGC TGC CTG TCT GTG ACC CAG AAA CCC ATC CCT GGG TAC ATC 267  
 Glu Asp Cys Cys Leu Ser Val Thr Gln Lys Pro Ile Pro Gly Tyr Ile 40  
 20 GTG AGG AAC TTC CAC TAC CTT CTC ATC AAG GAT GGC TGC AGG GTG CCT 315  
 Val Arg Asn Phe His Tyr Leu Leu Ile Lys Asp Gly Cys Arg Val Pro 55  
 25 GCT GTA GTG TTC ACC ACA CTG AGG GGC CGC CAG CTC TGT GCA CCC CCA 363  
 Ala Val Val Phe Thr Thr Leu Arg Gly Arg Gln Leu Cys Ala Pro Pro 70  
 30 GAC CAG CCC TGG GTA GAA CGC ATC ATC CAG AGA CTG CAG AGG ACC TCA 411  
 Asp Gln Pro Trp Val Glu Arg Ile Ile Gln Arg Leu Gln Arg Thr Ser 90  
 35 GCC AAG ATG AAG CGC CGC AGC AGT TAACCTATGA CCGTGCAGAG GGAGCCCGBA 465  
 Ala Lys Met Lys Arg Arg Ser Ser 95  
 GTCCGAGTCA AGCATTGTGA ATTATTACCT AACCTGGGGA ACCGAGGACC AGAAGGAAGG 525  
 ACCAGGCTTC CAGCTCCTCT GCACCAGACC TGACCAGCCA GGACAGGGCC TGGGGTGTGT 585  
 40 GTGAGTGTGA GTGTGAGCGA GAGGGTGAGT GTGGTCTAGA GTAAAGCTGC TCCACCCCCA 645  
 GATTGCAATG CTACCAATAA AGCCGCCTGG TGTTTACAAC TAAAAA AAAA 699

45

## (2) INFORMATION FOR SEQ ID NO:8:

## (i) SEQUENCE CHARACTERISTICS:

- 50 (A) LENGTH: 98 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## 55 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Met Ala Leu Leu Leu Ala Leu Ser Leu Leu Val Leu Trp Thr Ser Pro  
 1 5 10 15  
 60 Ala Pro Thr Leu Ser Gly Thr Asn Asp Ala Glu Asp Cys Cys Leu Ser

	20	25	30
	Val Thr Gln Lys Pro Ile Pro Gly Tyr Ile Val Arg Asn Phe His Tyr		
	35	40	45
5	Leu Leu Ile Lys Asp Gly Cys Arg Val Pro Ala Val Val Phe Thr Thr		
	50	55	60
10	Leu Arg Gly Arg Gln Leu Cys Ala Pro Pro Asp Gln Pro Trp Val Glu		
	65	70	75
	Arg Ile Ile Gln Arg Leu Gln Arg Thr Ser Ala Lys Met Lys Arg Arg		
	85	90	95
15	Ser Ser		

## (2) INFORMATION FOR SEQ ID NO:9:

20 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 1119 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: cDNA

30 (ix) FEATURE:  
 (A) NAME/KEY: CDS  
 (B) LOCATION: 1..1095

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

35	ATG TTT TCG ACT CCA GTG AAG ATT ATT TTG TGT CAG TCA ATA CTT CAT	48
	Met Phe Ser Thr Pro Val Lys Ile Ile Leu Cys Gln Ser Ile Leu His	
	1 5 10 15	
40	ATT ACT CAG TTG ATT CTG AGA TGT TAC TGT GCT CCT TGC AGG AGG TCA	96
	Ile Thr Gln Leu Ile Leu Arg Cys Tyr Cys Ala Pro Cys Arg Arg Ser	
	20 25 30	
45	GGC AGT TCT CCA GGC TAT TTG TAC CGA ATT GCC TAC TCC TTG ATC TGT	144
	Gly Ser Ser Pro Gly Tyr Leu Tyr Arg Ile Ala Tyr Ser Leu Ile Cys	
	35 40 45	
50	GTT CTT GGC CTC CTG GGG AAT ATT CTG GTG GTG ATC ACC TTT GCT TTT	192
	Val Leu Gly Leu Leu Gly Asn Ile Leu Val Val Ile Thr Phe Ala Phe	
	50 55 60	
55	TAT AAG AAG GCC AGG TCT ATG ACA GAC GTC TAT CTC TTG AAC ATG GCC	240
	Tyr Lys Lys Ala Arg Ser Met Thr Asp Val Tyr Leu Leu Asn Met Ala	
	65 70 75 80	
60	ATT GCA GAC ATC CTC TTT GTT CTT ACT CTC CCA TTC TGG GCA GTG AGT	288
	Ile Ala Asp Ile Leu Phe Val Leu Thr Leu Pro Phe Trp Ala Val Ser	
	85 90 95	
60	CAT GCC ACT GGT GCG TGG GTT TTC AGC AAT GCC ACG TGC AAG TTG CTA	336

	His	Ala	Thr	Gly	Ala	Trp	Val	Phe	Ser	Asn	Ala	Thr	Cys	Lys	Leu	Leu	
				100					105					110			
5	AAA	GGC	ATC	TAT	GCC	ATC	AAC	TTT	AAC	TGC	GGG	ATG	CTG	CTC	CTG	ACT	384
	Lys	Gly	Ile	Tyr	Ala	Ile	Asn	Phe	Asn	Cys	Gly	Met	Leu	Leu	Leu	Thr	
			115					120					125				
10	TGC	ATT	AGC	ATG	GAC	CGG	TAC	ATC	GCC	ATT	GTA	CAG	GCG	ACT	AAG	TCA	432
	Cys	Ile	Ser	Met	Asp	Arg	Tyr	Ile	Ala	Ile	Val	Gln	Ala	Thr	Lys	Ser	
			130				135					140					
15	TTC	CGG	CTC	CGA	TCC	AGA	ACA	CTA	CCG	CGC	AGC	AAA	ATC	ATC	TGC	CTT	480
	Phe	Arg	Leu	Arg	Ser	Arg	Thr	Leu	Pro	Arg	Ser	Lys	Ile	Ile	Cys	Leu	
	145					150					155					160	
	GTT	GTG	TGG	GGG	CTG	TCA	GTC	ATC	ATC	TCC	AGC	TCA	ACT	TTT	GTC	TTC	528
	Val	Val	Trp	Gly	Leu	Ser	Val	Ile	Ile	Ser	Ser	Ser	Thr	Phe	Val	Phe	
					165					170					175		
20	AAC	CAA	AAA	TAC	AAC	ACC	CAA	GGC	AGC	GAT	GTC	TGT	GAA	CCC	AAG	TAC	576
	Asn	Gln	Lys	Tyr	Asn	Thr	Gln	Gly	Ser	Asp	Val	Cys	Glu	Pro	Lys	Tyr	
				180					185					190			
25	CAA	ACT	GTC	TCG	GAG	CCC	ATC	AGG	TGG	AAG	CTG	CTG	ATG	TTG	GGG	CTT	624
	Gln	Thr	Val	Ser	Glu	Pro	Ile	Arg	Trp	Lys	Leu	Leu	Met	Leu	Gly	Leu	
			195					200					205				
30	GAG	CTA	CTC	TTT	GGT	TTC	TTT	ATC	CCT	TTG	ATG	TTC	ATG	ATA	TTT	TGT	672
	Glu	Leu	Leu	Phe	Gly	Phe	Phe	Ile	Pro	Leu	Met	Phe	Met	Ile	Phe	Cys	
		210					215					220					
35	TAC	ACG	TTC	ATT	GTC	AAA	ACC	TTG	GTG	CAA	GCT	CAG	AAT	TCT	AAA	AGG	720
	Tyr	Thr	Phe	Ile	Val	Lys	Thr	Leu	Val	Gln	Ala	Gln	Asn	Ser	Lys	Arg	
	225					230				235						240	
	CAC	AAA	GCC	ATC	CGT	GTA	ATC	ATA	GCT	GTG	GTG	CTT	GTG	TTT	CTG	GCT	768
	His	Lys	Ala	Ile	Arg	Val	Ile	Ile	Ala	Val	Val	Leu	Val	Phe	Leu	Ala	
					245					250					255		
40	TGT	CAG	ATT	CCT	CAT	AAC	ATG	GTC	CTG	CTT	GTG	ACG	GCT	GCT	AAT	TTG	816
	Cys	Gln	Ile	Pro	His	Asn	Met	Val	Leu	Val	Thr	Ala	Ala	Asn	Leu		
				260				265						270			
45	GGT	AAA	ATG	AAC	CGA	TCC	TGC	CAG	AGC	GAA	AAG	CTA	ATT	GGC	TAT	ACG	864
	Gly	Lys	Met	Asn	Arg	Ser	Cys	Gln	Ser	Glu	Lys	Leu	Ile	Gly	Tyr	Thr	
			275					280					285				
50	AAA	ACT	GTC	ACA	GAA	GTC	CTG	GCT	TTC	CTG	CAC	TGC	TGC	CTG	AAC	CCT	912
	Lys	Thr	Val	Thr	Glu	Val	Leu	Ala	Phe	Leu	His	Cys	Cys	Leu	Asn	Pro	
			290				295					300					
55	GTG	CTC	TAC	GCT	TTT	ATT	GGG	CAG	AAG	TTC	AGA	AAC	TAC	TTT	CTG	AAG	960
	Val	Leu	Tyr	Ala	Phe	Ile	Gly	Gln	Lys	Phe	Arg	Asn	Tyr	Phe	Leu	Lys	
	305					310				315						320	
	ATC	TTG	AAG	GAC	CTG	TGG	TGT	GTG	AGA	AGG	AAG	TAC	AAG	TCC	TCA	GGC	1008
	Ile	Leu	Lys	Asp	Leu	Trp	Cys	Val	Arg	Arg	Lys	Tyr	Lys	Ser	Ser	Gly	
					325					330					335		
60	TTC	TCC	TGT	GCC	GGG	AGG	TAC	TCA	GAA	AAC	ATT	TCT	CGG	CAG	ACC	AGT	1056

Phe Ser Cys Ala Gly Arg Tyr Ser Glu Asn Ile Ser Arg Gln Thr Ser  
 340 345 350

5 GAG ACC GCA GAT AAC GAC AAT GCG TCG TCC TTC ACT ATG TGATAGAAAG 1105  
 Glu Thr Ala Asp Asn Asp Asn Ala Ser Ser Phe Thr Met  
 355 360 365

CTGAGTCTCC CTAA

1119

10

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

15 (A) LENGTH: 365 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Met Phe Ser Thr Pro Val Lys Ile Ile Leu Cys Gln Ser Ile Leu His  
 1 5 10 15  
 25 Ile Thr Gln Leu Ile Leu Arg Cys Tyr Cys Ala Pro Cys Arg Arg Ser  
 20 25 30  
 Gly Ser Ser Pro Gly Tyr Leu Tyr Arg Ile Ala Tyr Ser Leu Ile Cys  
 35 40 45  
 30 Val Leu Gly Leu Leu Gly Asn Ile Leu Val Val Ile Thr Phe Ala Phe  
 50 55 60  
 Tyr Lys Lys Ala Arg Ser Met Thr Asp Val Tyr Leu Leu Asn Met Ala  
 65 70 75 80  
 35 Ile Ala Asp Ile Leu Phe Val Leu Thr Leu Pro Phe Trp Ala Val Ser  
 85 90 95  
 40 His Ala Thr Gly Ala Trp Val Phe Ser Asn Ala Thr Cys Lys Leu Leu  
 100 105 110  
 Lys Gly Ile Tyr Ala Ile Asn Phe Asn Cys Gly Met Leu Leu Leu Thr  
 115 120 125  
 45 Cys Ile Ser Met Asp Arg Tyr Ile Ala Ile Val Gln Ala Thr Lys Ser  
 130 135 140  
 Phe Arg Leu Arg Ser Arg Thr Leu Pro Arg Ser Lys Ile Ile Cys Leu  
 145 150 155 160  
 50 Val Val Trp Gly Leu Ser Val Ile Ile Ser Ser Ser Thr Phe Val Phe  
 165 170 175  
 55 Asn Gln Lys Tyr Asn Thr Gln Gly Ser Asp Val Cys Glu Pro Lys Tyr  
 180 185 190  
 Gln Thr Val Ser Glu Pro Ile Arg Trp Lys Leu Leu Met Leu Gly Leu  
 195 200 205

60

Glu Leu Leu Phe Gly Phe Phe Ile Pro Leu Met Phe Met Ile Phe Cys  
 210 215 220  
 Tyr Thr Phe Ile Val Lys Thr Leu Val Gln Ala Gln Asn Ser Lys Arg  
 5 225 230 235 240  
 His Lys Ala Ile Arg Val Ile Ile Ala Val Val Leu Val Phe Leu Ala  
 245 250 255  
 10 Cys Gln Ile Pro His Asn Met Val Leu Leu Val Thr Ala Ala Asn Leu  
 260 265 270  
 Gly Lys Met Asn Arg Ser Cys Gln Ser Glu Lys Leu Ile Gly Tyr Thr  
 275 280 285  
 15 Lys Thr Val Thr Glu Val Leu Ala Phe Leu His Cys Cys Leu Asn Pro  
 290 295 300  
 Val Leu Tyr Ala Phe Ile Gly Gln Lys Phe Arg Asn Tyr Phe Leu Lys  
 20 305 310 315 320  
 Ile Leu Lys Asp Leu Trp Cys Val Arg Arg Lys Tyr Lys Ser Ser Gly  
 325 330 335  
 25 Phe Ser Cys Ala Gly Arg Tyr Ser Glu Asn Ile Ser Arg Gln Thr Ser  
 340 345 350  
 Glu Thr Ala Asp Asn Asp Asn Ala Ser Ser Phe Thr Met  
 355 360 365  
 30 (2) INFORMATION FOR SEQ ID NO:11:  
 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 1547 base pairs  
 35 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear  
 (ii) MOLECULE TYPE: cDNA  
 40  
 (ix) FEATURE:  
 (A) NAME/KEY: CDS  
 (B) LOCATION: 49..1116  
 45  
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:  
 50 GAGGAAGCTG CTTCTGGGGGG TGAGCAAACT TTTTAAAATG CAGAAATT ATG ATC TAC 57  
 Met Ile Tyr  
 1  
 ACC CGT TTC TTA AAA GGC AGT CTG AAG ATG GCC AAT TAC ACG CTG GCA 105  
 Thr Arg Phe Leu Lys Gly Ser Leu Lys Met Ala Asn Tyr Thr Leu Ala  
 55 5 10 15  
 CCA GAG GAT GAA TAT GAT GTC CTC ATA GAA GGT GAA CTG GAG AGC GAT 153  
 Pro Glu Asp Glu Tyr Asp Val Leu Ile Glu Gly Glu Leu Glu Ser Asp  
 20 25 30 35  
 60



	GAG GCA GAG CAA TGT GAC AAG TAT GAC GCC CAG GCA CTC TCA GCC CAG	201
	Glu Ala Glu Gln Cys Asp Lys Tyr Asp Ala Gln Ala Leu Ser Ala Gln	
	40 45 50	
5	CTG GTG CCA TCA CTC TGC TCT GCT GTG TTT GTG ATC GGT GTC CTG GAC	249
	Leu Val Pro Ser Leu Cys Ser Ala Val Phe Val Ile Gly Val Leu Asp	
	55 60 65	
10	AAT CTC CTG GTT GTG CTT ATC CTG GTA AAA TAT AAA GGA CTC AAA CGC	297
	Asn Leu Leu Val Val Leu Ile Leu Val Lys Tyr Lys Gly Leu Lys Arg	
	70 75 80	
15	GTG GAA AAT ATC TAT CTT CTA AAC TTG GCA GTT TCT AAC TTG TGT TTC	345
	Val Glu Asn Ile Tyr Leu Leu Asn Leu Ala Val Ser Asn Leu Cys Phe	
	85 90 95	
20	TTG CTT ACC CTG CCC TTC TGG GCT CAT GCT GGG GGC GAT CCC ATG TGT	393
	Leu Leu Thr Leu Pro Phe Trp Ala His Ala Gly Gly Asp Pro Met Cys	
	100 105 110 115	
20	AAA ATT CTC ATT GGA CTG TAC TTC GTG GGC CTG TAC AGT GAG ACA TTT	441
	Lys Ile Leu Ile Gly Leu Tyr Phe Val Gly Leu Tyr Ser Glu Thr Phe	
	120 125 130	
25	TTC AAT TGC CTT CTG ACT GTG CAA AGG TAC CTA GTG TTT TTG CAC AAG	489
	Phe Asn Cys Leu Leu Thr Val Gln Arg Tyr Leu Val Phe Leu His Lys	
	135 140 145	
30	GGC AAC TTT TTC TCA GCC AGG AGG AGG GTG CCC TGT GGC ATC ATT ACA	537
	Gly Asn Phe Phe Ser Ala Arg Arg Arg Val Pro Cys Gly Ile Ile Thr	
	150 155 160	
35	AGT GTC CTG GCA TGG GTA ACA GCC ATT CTG GCC ACT TTG CCT GAA TTC	585
	Ser Val Leu Ala Trp Val Thr Ala Ile Leu Ala Thr Leu Pro Glu Phe	
	165 170 175	
40	GTG GTT TAT AAA CCT CAG ATG GAA GAC CAG AAA TAC AAG TGT GCA TTT	633
	Val Val Tyr Lys Pro Gln Met Glu Asp Gln Lys Tyr Lys Cys Ala Phe	
	180 185 190 195	
40	AGC AGA ACT CCC TTC CTG CCA GCT GAT GAG ACA TTC TGG AAG CAT TTT	681
	Ser Arg Thr Pro Phe Leu Pro Ala Asp Glu Thr Phe Trp Lys His Phe	
	200 205 210	
45	CTG ACT TTA AAA ATG AAC ATT TCG GTT CTT GTC CTC CCC CTA TTT ATT	729
	Leu Thr Leu Lys Met Asn Ile Ser Val Leu Val Leu Pro Leu Phe Ile	
	215 220 225	
50	TTT ACA TTT CTC TAT GTG CAA ATG AGA AAA ACA CTA AGG TTC AGG GAG	777
	Phe Thr Phe Leu Tyr Val Gln Met Arg Lys Thr Leu Arg Phe Arg Glu	
	230 235 240	
55	CAG AGG TAT AGC CTT TTC AAG CTT GTT TTT GCC GTA ATG GTA GTC TTC	825
	Gln Arg Tyr Ser Leu Phe Lys Leu Val Phe Ala Val Met Val Val Phe	
	245 250 255	
60	CTT CTG ATG TGG GCG CCC TAC AAT ATT GCA TTT TTC CTG TCC ACT TTC	873
	Leu Leu Met Trp Ala Pro Tyr Asn Ile Ala Phe Phe Leu Ser Thr Phe	
	260 265 270 275	

	AAA GAA CAC TTC TCC CTG AGT GAC TGC AAG AGC AGC TAC AAT CTG GAC	921
	Lys Glu His Phe Ser Leu Ser Asp Cys Lys Ser Ser Tyr Asn Leu Asp	
	280 285 290	
5	AAA AGT GTT CAC ATC ACT AAA CTC ATC GCC ACC ACC CAC TGC TGC ATC	969
	Lys Ser Val His Ile Thr Lys Leu Ile Ala Thr Thr His Cys Cys Ile	
	295 300 305	
10	AAC CCT CTC CTG TAT GCG TTT CTT GAT GGG ACA TTT AGC AAA TAC CTC	1017
	Asn Pro Leu Leu Tyr Ala Phe Leu Asp Gly Thr Phe Ser Lys Tyr Leu	
	310 315 320	
15	TGC CGC TGT TTC CAT CTG CGT AGT AAC ACC CCA CTT CAA CCC AGG GGG	1065
	Cys Arg Cys Phe His Leu Arg Ser Asn Thr Pro Leu Gln Pro Arg Gly	
	325 330 335	
20	CAG TCT GCA CAA GGC ACA TCG AGG GAA GAA CCT GAC CAT TCC ACC GAA	1113
	Gln Ser Ala Gln Gly Thr Ser Arg Glu Glu Pro Asp His Ser Thr Glu	
	340 345 350 355	
25	GTG TAAACTAGCA TCCACCAAAT GCAAGAAGAA TAAACATGGA TTTTCATCTT	1166
	Val	
30	TCTGCATTAT TTCATGTAAA TTTTCTACAC ATTTGTATAC AAAATCGGAT ACAGGAAGAA	1226
	AAGGGAGAGG TGAGCTAACA TTTGCTAAGC ACTGAATTTG TCTCAGGCAC CGTGCAAGGC	1286
	TCTTTACAAA CGTGAGCTCC TTCGCCTCCT ACCACTTGTC CATAGTGTGG ATAGGACTAG	1346
35	TCTCATTTCT CTGAGAAGAA AACTAAGGCG CGGAAATTTG TCTAAGATCA CATAACTAGG	1406
	AAGTGGCAGA ACTGATTCTC CAGCCCTGGT AGCATTGCT CAGAGCCTAC GCTTGGTCCA	1466
	GAACATCAAAA CTCCAAACCC TGGGGACAAA CGACATGAAA TAAATGTATT TTAAAACATA	1526
	TAAAAAAAAA AAAAAAAAAA A	1547
40	(2) INFORMATION FOR SEQ ID NO:12:	
	(i) SEQUENCE CHARACTERISTICS:	
	(A) LENGTH: 356 amino acids	
	(B) TYPE: amino acid	
45	(D) TOPOLOGY: linear	
	(ii) MOLECULE TYPE: protein	
	(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:	
50	Met Ile Tyr Thr Arg Phe Leu Lys Gly Ser Leu Lys Met Ala Asn Tyr	
	1 5 10 15	
55	Thr Leu Ala Pro Glu Asp Glu Tyr Asp Val Leu Ile Glu Gly Glu Leu	
	20 25 30	
	Glu Ser Asp Glu Ala Glu Gln Cys Asp Lys Tyr Asp Ala Gln Ala Leu	
	35 40 45	
60	Ser Ala Gln Leu Val Pro Ser Leu Cys Ser Ala Val Phe Val Ile Gly	

50 55 60

Val Leu Asp Asn Leu Leu Val Val Leu Ile Leu Val Lys Tyr Lys Gly  
65 70 75 80

5 Leu Lys Arg Val Glu Asn Ile Tyr Leu Leu Asn Leu Ala Val Ser Asn  
85 90 95

10 Leu Cys Phe Leu Leu Thr Leu Pro Phe Trp Ala His Ala Gly Gly Asp  
100 105 110

Pro Met Cys Lys Ile Leu Ile Gly Leu Tyr Phe Val Gly Leu Tyr Ser  
115 120 125

15 Glu Thr Phe Phe Asn Cys Leu Leu Thr Val Gln Arg Tyr Leu Val Phe  
130 135 140

Leu His Lys Gly Asn Phe Phe Ser Ala Arg Arg Arg Val Pro Cys Gly  
145 150 155 160

20 Ile Ile Thr Ser Val Leu Ala Trp Val Thr Ala Ile Leu Ala Thr Leu  
165 170 175

Pro Glu Phe Val Val Tyr Lys Pro Gln Met Glu Asp Gln Lys Tyr Lys  
180 185 190

25 Cys Ala Phe Ser Arg Thr Pro Phe Leu Pro Ala Asp Glu Thr Phe Trp  
195 200 205

30 Lys His Phe Leu Thr Leu Lys Met Asn Ile Ser Val Leu Val Leu Pro  
210 215 220

Leu Phe Ile Phe Thr Phe Leu Tyr Val Gln Met Arg Lys Thr Leu Arg  
225 230 235 240

35 Phe Arg Glu Gln Arg Tyr Ser Leu Phe Lys Leu Val Phe Ala Val Met  
245 250 255

Val Val Phe Leu Leu Met Trp Ala Pro Tyr Asn Ile Ala Phe Phe Leu  
260 265 270

40 Ser Thr Phe Lys Glu His Phe Ser Leu Ser Asp Cys Lys Ser Ser Tyr  
275 280 285

45 Asn Leu Asp Lys Ser Val His Ile Thr Lys Leu Ile Ala Thr Thr His  
290 295 300

Cys Cys Ile Asn Pro Leu Leu Tyr Ala Phe Leu Asp Gly Thr Phe Ser  
305 310 315 320

50 Lys Tyr Leu Cys Arg Cys Phe His Leu Arg Ser Asn Thr Pro Leu Gln  
325 330 335

Pro Arg Gly Gln Ser Ala Gln Gly Thr Ser Arg Glu Glu Pro Asp His  
340 345 350

55 Ser Thr Glu Val  
355

60 (2) INFORMATION FOR SEQ ID NO:13:

## (i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 355 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

10

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

15 Met Glu Thr Pro Asn Thr Thr Glu Asp Tyr Asp Thr Thr Thr Glu Phe  
 1 5 10 15

Asp Tyr Gly Asp Ala Thr Pro Cys Gln Lys Val Asn Glu Arg Ala Phe  
 20 20 25 30

Gly Ala Gln Leu Leu Pro Pro Leu Tyr Ser Leu Val Phe Val Ile Gly  
 35 40 45

25 Leu Val Gly Asn Ile Leu Val Val Leu Val Leu Val Gln Tyr Lys Arg  
 50 55 60

Leu Lys Asn Met Thr Ser Ile Tyr Leu Leu Asn Leu Ala Ile Ser Asp  
 65 70 75 80

30 Leu Leu Phe Leu Phe Thr Leu Pro Phe Trp Ile Asp Tyr Lys Leu Lys  
 85 90 95

Asp Asp Trp Val Phe Gly Asp Ala Met Cys Lys Ile Leu Ser Gly Phe  
 100 105 110

35 Tyr Tyr Thr Gly Leu Tyr Ser Glu Ile Phe Phe Ile Ile Leu Leu Thr  
 115 120 125

Ile Asp Arg Tyr Leu Ala Ile Val His Ala Val Phe Ala Leu Arg Ala  
 40 130 135 140

Arg Thr Val Thr Phe Gly Val Ile Thr Ser Ile Ile Ile Trp Ala Leu  
 145 150 155 160

45 Ala Ile Leu Ala Ser Met Pro Gly Leu Tyr Phe Ser Lys Thr Gln Trp  
 165 170 175

Glu Phe Thr His His Thr Cys Ser Leu His Phe Pro His Glu Ser Leu  
 180 185 190

50 Arg Glu Trp Lys Leu Phe Gln Ala Leu Lys Leu Asn Leu Phe Gly Leu  
 195 200 205

Val Leu Pro Leu Leu Val Met Ile Ile Cys Tyr Thr Gly Ile Ile Lys  
 55 210 215 220

Ile Leu Leu Arg Arg Pro Asn Glu Lys Lys Ser Lys Ala Val Arg Leu  
 225 230 235 240

60 Ile Phe Val Ile Met Ile Ile Phe Phe Leu Phe Trp Thr Pro Tyr Asn

245 250 255

Leu Thr Ile Leu Ile Ser Val Phe Gln Asp Phe Leu Phe Thr His Glu  
260 265 270

5 Cys Glu Gln Ser Arg His Leu Asp Leu Ala Val Gln Val Thr Glu Val  
275 280 285

Ile Ala Tyr Thr His Cys Cys Val Asn Pro Val Ile Tyr Ala Phe Val  
290 295 300

10 Gly Glu Arg Phe Arg Lys Tyr Leu Arg Gln Leu Phe His Arg Arg Val  
305 310 315 320

15 Ala Val His Leu Val Lys Trp Leu Pro Phe Leu Ser Val Asp Arg Leu  
325 330 335

Glu Arg Val Ser Ser Thr Ser Pro Ser Thr Gly Glu His Glu Leu Ser  
340 345 350

20 Ala Gly Phe  
355

## (2) INFORMATION FOR SEQ ID NO:14:

25

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 374 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

30

- (ii) MOLECULE TYPE: protein

35

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

40

Met Leu Ser Thr Ser Arg Ser Arg Phe Ile Arg Asn Thr Asn Glu Ser  
1 5 10 15

Gly Glu Glu Val Thr Thr Phe Phe Asp Tyr Asp Tyr Gly Ala Pro Cys  
20 25 30

45

His Lys Phe Asp Val Lys Gln Ile Gly Ala Gln Leu Leu Pro Pro Leu  
35 40 45

Tyr Ser Leu Val Phe Ile Phe Gly Phe Val Gly Asn Met Leu Val Val  
50 55 60

50

Leu Ile Leu Ile Asn Cys Lys Lys Leu Lys Cys Leu Thr Asp Ile Tyr  
65 70 75 80

55

Leu Leu Asn Leu Ala Ile Ser Asp Leu Leu Phe Leu Ile Thr Leu Pro  
85 90 95

Leu Trp Ala His Ser Ala Ala Asn Glu Trp Val Phe Gly Asn Ala Met  
100 105 110

60

Cys Lys Leu Phe Thr Gly Leu Tyr His Ile Gly Tyr Phe Gly Gly Ile

	115	120	125
	Phe Phe Ile Ile Leu Leu Thr Ile Asp Arg Tyr Leu Ala Ile Val His		
	130	135	140
5	Ala Val Phe Ala Leu Lys Ala Arg Thr Val Thr Phe Gly Val Val Thr		
	145	150	155 160
10	Ser Val Ile Thr Trp Leu Val Ala Val Phe Ala Ser Val Pro Gly Ile		
		165	170 175
	Ile Phe Thr Lys Cys Gln Lys Glu Asp Ser Val Tyr Val Cys Gly Pro		
		180	185 190
15	Tyr Phe Pro Arg Gly Trp Asn Asn Phe His Thr Ile Met Arg Asn Ile		
		195	200 205
	Leu Gly Leu Val Leu Pro Leu Leu Ile Met Val Ile Cys Tyr Ser Gly		
		210	215 220
20	Ile Leu Lys Thr Leu Leu Arg Cys Arg Asn Glu Lys Lys Arg His Arg		
		225	230 235 240
	Ala Val Arg Val Ile Phe Thr Ile Met Ile Val Tyr Phe Leu Phe Trp		
		245	250 255
25	Thr Pro Tyr Asn Ile Val Ile Leu Leu Asn Thr Phe Gln Glu Phe Phe		
		260	265 270
30	Gly Leu Ser Asn Cys Glu Ser Thr Ser Gln Leu Asp Gln Ala Thr Gln		
		275	280 285
	Val Thr Glu Thr Leu Gly Met Thr His Cys Cys Ile Asn Pro Ile Ile		
		290	295 300
35	Tyr Ala Phe Val Gly Glu Lys Phe Arg Ser Leu Phe His Ile Ala Leu		
		305	310 315 320
	Gly Cys Arg Ile Ala Pro Leu Gln Lys Pro Val Cys Gly Gly Pro Gly		
		325	330 335
40	Val Arg Pro Gly Lys Asn Val Lys Val Thr Thr Gln Gly Leu Leu Asp		
		340	345 350
45	Gly Arg Gly Lys Gly Lys Ser Ile Gly Arg Ala Pro Glu Ala Ser Leu		
		355	360 365
	Gln Asp Lys Glu Gly Ala		
		370	

## (2) INFORMATION FOR SEQ ID NO:15:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 355 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

5 Met Thr Thr Ser Leu Asp Thr Val Glu Thr Phe Gly Thr Thr Ser Tyr  
 1 5 10 15  
 10 Tyr Asp Asp Val Gly Leu Leu Cys Glu Lys Ala Asp Thr Arg Ala Leu  
 20 25 30  
 Met Ala Gln Phe Val Pro Pro Leu Tyr Ser Leu Val Phe Thr Val Gly  
 35 40 45  
 15 Leu Leu Gly Asn Val Val Val Val Met Ile Leu Ile Lys Tyr Arg Arg  
 50 55 60  
 Leu Arg Ile Met Thr Asn Ile Tyr Leu Leu Asn Leu Ala Ile Ser Asp  
 65 70 75 80  
 20 Leu Leu Phe Leu Val Thr Leu Pro Phe Trp Ile His Tyr Val Arg Gly  
 85 90 95  
 His Asn Trp Val Phe Gly His Gly Met Cys Lys Leu Leu Ser Gly Phe  
 100 105 110  
 25 Tyr His Thr Gly Leu Tyr Ser Glu Ile Phe Phe Ile Ile Leu Leu Thr  
 115 120 125  
 30 Ile Asp Arg Tyr Leu Ala Ile Val His Ala Val Phe Ala Leu Arg Ala  
 130 135 140  
 Arg Thr Val Thr Phe Gly Val Ile Thr Ser Ile Val Thr Trp Gly Leu  
 145 150 155 160  
 35 Ala Val Leu Ala Ala Leu Pro Glu Phe Ile Phe Tyr Glu Thr Glu Glu  
 165 170 175  
 Leu Phe Glu Glu Thr Leu Cys Ser Ala Leu Tyr Pro Glu Asp Thr Val  
 180 185 190  
 40 Tyr Ser Trp Arg His Phe His Thr Leu Arg Met Thr Ile Phe Cys Leu  
 195 200 205  
 45 Val Leu Pro Leu Leu Val Met Ala Ile Cys Tyr Thr Gly Ile Ile Lys  
 210 215 220  
 Thr Leu Leu Arg Cys Pro Ser Lys Lys Lys Tyr Lys Ala Ile Arg Leu  
 225 230 235 240  
 50 Ile Phe Val Ile Met Ala Val Phe Phe Ile Phe Trp Thr Pro Tyr Asn  
 245 250 255  
 Val Ala Ile Leu Leu Ser Ser Tyr Gln Ser Ile Leu Phe Gly Asn Asp  
 260 265 270  
 55 Cys Glu Arg Ser Lys His Leu Asp Leu Val Met Leu Val Thr Glu Val  
 275 280 285  
 60 Ile Ala Tyr Ser His Cys Cys Met Asn Pro Val Ile Tyr Ala Phe Val

290 295 300

Gly Glu Arg Phe Arg Lys Tyr Leu Arg His Phe Phe His Arg His Leu  
305 310 315 320

5 Leu Met His Leu Gly Arg Tyr Ile Pro Phe Leu Pro Ser Glu Lys Leu  
325 330 335

10 Glu Arg Thr Ser Ser Val Ser Pro Ser Thr Ala Glu Pro Glu Leu Ser  
340 345 350

Ile Val Phe  
355

15 (2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 360 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

20 (ii) MOLECULE TYPE: protein

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

30 Met Asn Pro Thr Asp Ile Ala Asp Thr Thr Leu Asp Glu Ser Ile Tyr  
1 5 10 15

Ser Asn Tyr Tyr Leu Tyr Glu Ser Ile Pro Lys Pro Cys Thr Lys Glu  
20 25 30

35 Gly Ile Lys Ala Phe Gly Glu Leu Phe Leu Pro Pro Leu Tyr Ser Leu  
35 40 45

40 Val Phe Val Phe Gly Leu Leu Gly Asn Ser Val Val Val Leu Val Leu  
50 55 60

Phe Lys Tyr Lys Arg Leu Arg Ser Met Thr Asp Val Tyr Leu Leu Asn  
65 70 75 80

45 Leu Ala Ile Ser Asp Leu Leu Phe Val Phe Ser Leu Pro Phe Trp Gly  
85 90 95

Tyr Tyr Ala Ala Asp Gln Trp Val Phe Gly Leu Gly Leu Cys Lys Met  
100 105 110

50 Ile Ser Trp Met Tyr Leu Val Gly Phe Tyr Ser Gly Ile Phe Phe Val  
115 120 125

55 Met Leu Met Ser Ile Asp Arg Tyr Leu Ala Ile Val His Ala Val Phe  
130 135 140

Ser Leu Arg Ala Arg Thr Leu Thr Tyr Gly Val Ile Thr Ser Leu Ala  
145 150 155 160

60 Thr Trp Ser Val Ala Val Phe Ala Ser Leu Pro Gly Phe Leu Phe Ser



165 170 175

Thr Cys Tyr Thr Glu Arg Asn His Thr Tyr Cys Lys Thr Lys Tyr Ser  
180 185 190

5 Leu Asn Ser Thr Thr Trp Lys Val Leu Ser Ser Leu Glu Ile Asn Ile  
195 200 205

10 Leu Gly Leu Val Ile Pro Leu Gly Ile Met Leu Phe Cys Tyr Ser Met  
210 215 220

Ile Ile Arg Thr Leu Gln His Cys Lys Asn Glu Lys Lys Asn Lys Ala  
225 230 235 240

15 Val Lys Met Ile Phe Ala Val Val Val Leu Phe Leu Gly Phe Trp Thr  
245 250 255

Pro Tyr Asn Ile Val Leu Phe Leu Glu Thr Leu Val Glu Leu Glu Val  
260 265 270

20 Leu Gln Asp Cys Thr Phe Glu Arg Tyr Leu Asp Tyr Ala Ile Gln Ala  
275 280 285

25 Thr Glu Thr Leu Ala Phe Val His Cys Cys Leu Asn Pro Ile Ile Tyr  
290 295 300

Phe Phe Leu Gly Glu Lys Phe Arg Lys Tyr Ile Leu Gln Leu Phe Lys  
305 310 315 320

30 Thr Cys Arg Gly Leu Phe Val Leu Cys Gln Tyr Cys Gly Leu Leu Gln  
325 330 335

Ile Tyr Ser Ala Asp Thr Pro Ser Ser Ser Tyr Thr Gln Ser Thr Met  
340 345 350

35 Asp His Asp Leu His Asp Ala Leu  
355 360

## (2) INFORMATION FOR SEQ ID NO:17:

- 40 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 23 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 45 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

50

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

GTAATGATCA GTCAACGGGG GAC

23

55

## (2) INFORMATION FOR SEQ ID NO:18:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 24 base pairs  
 60 (B) TYPE: nucleic acid

(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

10

CCAGCAAGCT TGCAACCTTA ACCA

24

(2) INFORMATION FOR SEQ ID NO:19:

15

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 9 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

20

(ii) MOLECULE TYPE: peptide

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

Asp Tyr Lys Asp Asp Asp Asp Lys Leu  
1 5

30

CCAGCAAGCTTGCAACCTTAACCA

## WHAT IS CLAIMED IS:

1. A substantially pure or isolated polypeptide comprising a segment exhibiting sequence homology to a corresponding portion of a mature protein selected from the group consisting of:
- i) TECK;
  - ii) MIP-3 $\alpha$ ;
  - iii) MIP-3 $\beta$ ;
  - iv) DC CR; and
  - v) M/DC CR;
- wherein said homology is at least about 70% identity and said portion is at least about 25 amino acids.
2. The protein of Claim 1, further comprising a second segment exhibiting:
- a) at least about 90% identity over at least 9 amino acids; or
  - b) at least about 80% identity over at least 17 amino acids.
3. The polypeptide of Claim 1, wherein said polypeptide:
- a) is from a warm blooded animal selected from the group of birds and mammals, including a mouse or human;
  - b) comprises a natural sequence from Tables 1 through 5;
  - c) exhibits a post-translational modification pattern distinct from a natural form of said polypeptide;
  - d) is made by expression of a recombinant nucleic acid;
  - e) comprises synthetic sequence;
  - f) is detectably labeled;
  - g) is conjugated to a solid substrate;
  - h) is conjugated to another chemical moiety;
  - i) is a fusion protein;

- j) is in a denatured conformation, including detergent denaturation;
- k) further comprises an epitope tag;
- l) is an immunogenic polypeptide;
- 5 m) has a defined homogeneous molecular weight;
- n) is useful as a carbon source;
- o) is an allelic variant of SEQ ID NO: 2, 4, 6, 8, 10, or 12;
- 10 p) is a 3-fold or less substituted form of a natural sequence;
- q) is in a sterile composition;
- r) is in a buffered solution or suspension;
- s) is in a regulated release device;
- t) comprises a post-translational modification;
- 15 u) is in a cell; or
- v) is in a kit which further comprises instructions for use or disposal of reagents therein.
4. An isolated or recombinant nucleic acid encoding said protein of Claim 1, where said portion consists of sequence from the coding region of SEQ ID NO: 1, 3, 5, 7, 9, or 11.
- 20
5. The nucleic acid of Claim 4, wherein said nucleic acid:
- 25
- a) exhibits at least about 80% identity to a natural cDNA encoding said segment;
- b) is in an expression vector;
- c) further comprises a promoter;
- 30 d) further comprises an origin of replication;
- e) is from a natural source;
- f) is detectably labeled;
- g) comprises synthetic nucleotide sequence;
- h) is less than 6 kb;
- 35 i) is from a mammal;
- j) comprises a natural full length mature coding sequence;

- k) is in a kit, which also comprises instructions for use or disposal of reagents therein;
- l) is a specific hybridization probe for a gene encoding said protein;
- 5 m) is a PCR product; or
- n) is in a cell.

6. A method of using a purified nucleic acid of Claim 5, comprising a step of expressing said nucleic acid to produce a protein.

10

7. An isolated or recombinant nucleic acid which encodes at least eight consecutive residues of SEQ ID NO: 2, 4, 6, 8, 10, or 12.

15

8. The nucleic acid of Claim 7, which encodes at least:

- a) twelve consecutive residues from SEQ ID NO: 2, and further comprises a coding sequence of at least 17 nucleotides from SEQ ID NO: 1;
- 20 b) twelve consecutive residues from SEQ ID NO: 4, and further comprises a coding sequence of at least 17 nucleotides from SEQ ID NO: 3;
- c) twelve consecutive residues from SEQ ID NO: 6, and further comprises a coding sequence of at least 17 nucleotides from SEQ ID NO: 5;
- 25 d) twelve consecutive residues from SEQ ID NO: 8, and further comprises a coding sequence of at least 17 nucleotides from SEQ ID NO: 7;
- e) twelve consecutive residues from SEQ ID NO: 10, and further comprises a coding sequence of at least 17 nucleotides from SEQ ID NO: 9; or
- 30 f) twelve consecutive residues from SEQ ID NO: 12, and further comprises a coding sequence of at least 17 nucleotides from SEQ ID NO: 11.
- 35

9. The nucleic acid of Claim 7, wherein said nucleic acid:

- a) exhibits at least about 80% identity to a natural cDNA encoding said segment;
- 5 b) is in an expression vector;
- c) further comprises a promoter;
- d) further comprises an origin of replication;
- e) encodes a 3-fold or less substituted sequence from a natural sequence;
- 10 f) is from a natural source;
- g) is detectably labeled;
- h) comprises synthetic nucleotide sequence;
- i) is less than 6 kb;
- j) is from a mammal;
- 15 k) is attached to a solid substrate, including in a Southern or Northern blot;
- l) comprises a natural full length coding sequence;
- m) is in a cell; or
- n) is in a detection kit, which also comprises  
20 instructions for use or disposal of reagents therein.

10. A nucleic acid which hybridizes under stringent wash conditions of 55° C and less than 150 mM salt to the  
25 nucleic acid of Claim 7.

11. The nucleic acid of Claim 10, which exhibits at least about 85% identity over a stretch of at least about 30 nucleotides to a primate sequence of SEQ ID NO: 1, 3, 5,  
30 7, 9, or 11.

12. The nucleic acid of Claim 10, wherein:  
a) said identity is at least 90%; or  
b) said stretch is at least 75 nucleotides.

35

13. The nucleic acid of Claim 10, wherein:  
a) said identity is at least 95%; or

b) said stretch is at least 100 nucleotides.

14. A binding compound comprising an antigen binding fragment from an antibody which binds to a protein of Claim

5 1.

15. The binding compound of Claim 14, wherein:

a) said polypeptide is a mouse or human protein;

10 b) said antibody is raised against a mature peptide sequence of Tables 1 through 5;

c) said antibody is a monoclonal antibody;

d) said binding compound is attached to a solid substrate;

e) said binding compound is in a sterile composition;

15 f) said binding compound binds to a denatured antigen, including a detergent denatured antigen;

g) said binding compound is detectably labeled;

h) said binding compound is an Fv, Fab, or Fab2 fragment;

20 i) said binding compound is conjugated to a chemical moiety;

j) said binding compound is in a detection kit which also comprises instructions for use or disposal of reagents therein.

25

16. A cell which makes said antibody of Claim 14.

17. A method of purifying a polypeptide using a binding compound of Claim 14 to specifically separate said  
30 polypeptides from others.

18. A method of generating an antigen-binding compound complex comprising the step of contacting a sample comprising said antigen to a sample comprising a binding  
35 compound of Claim 14.

19. A method of modulating physiology or development of a cell expressing a receptor for a chemokine selected from the group selected from:

- a) TECK;
- 5 b) MIP-3 $\alpha$ ; or
- c) MIP-3 $\beta$ ;

comprising contacting said cell with a composition comprising:

- i) an agonist or mutein of said chemokine; or
- 10 ii) an antibody antagonist of said chemokine.

20. The method of Claim 19, wherein said cell is a macrophage or lymphocyte.

15 21. The method of Claim 19, wherein said physiology is selected from:

- a) a cellular calcium flux;
- b) a chemoattractant response;
- c) cellular morphology modification responses;
- 20 d) phosphoinositide lipid turnover; or
- e) an antiviral response.

22. The method of Claim 19, wherein:

- a) said receptor is DC CR and said chemokine is MIP-3 $\alpha$ ;
- 25 b) said physiology is pulmonary physiology; or
- c) said cell is an eosinophil.



## MAMMALIAN CHEMOKINE REAGENTS



5

## ABSTRACT

Novel chemokines from mammals, reagents related thereto including purified proteins, specific antibodies, and nucleic acids encoding said chemokines. Chemokine  
10 receptors are also provided. Methods of using said reagents and diagnostic kits are also provided.

This filing is a regular U.S. Patent Application  
15 resulting from the conversion of provisional applications DX0589P, USSN 60/021,664, filed July 5, 1996; DX0589P1, USSN 60/028,329, filed October 11, 1996; DX0589P2 (Wang, et al.) no Application Number yet assigned, filed June 4, 1997; each of which is incorporated herein by reference,  
20 and priority claimed to.

46225-462880

**1A****1B**

DECLARATION AND POWER OF  
ATTORNEY FOR PATENT APPLICATION

Attorney's Docket No. DX0589Q

As a below-named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name;

I believe I am the original, first sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

"MAMMALIAN CHEMOKINE REAGENTS"

the specification of which

☒ is attached hereto.

☐ was filed on July 3, 1997 as Application Serial No.                      to be assigned

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, §1.56(a).

I hereby claim foreign priority benefits under Title 35, United States Code, §119(a)-(d) of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

Prior Foreign Application(s):	Priority Claimed		
<u>                    </u> (Number)	<u>                    </u> (Country)	<u>                    </u> (Day/Month/Year Filed)	<u>                    </u> Yes or No

I hereby claim the benefit under Title 35, United States Code, §119(e) of any United States provisional application(s) listed below:

<u>60/021,664</u> (Application Number)	<u>July 5, 1996</u> (Filing Date)	<u>Kurt GISH, et al.</u> (Inventor(s))
<u>60/028,329</u> (Application Number)	<u>October 11, 1996</u> (Filing Date)	<u>Kurt GISH, et al.</u> (Inventor(s))
<u>to be assigned</u> (Application Number)	<u>June 4, 1997</u> (Filing Date)	<u>Wei WANG, et al.</u> (Inventor(s))

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose material information as

defined in Title 37, Code of Federal Regulations, §1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

(Application Serial No.)

(Filing Date)

(Status – patented, pending, abandoned)

Power of Attorney: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in Patent and Trademark Office connected therewith. (List name and registration number.)

Edwin P. Ching	Reg. No. 34,090	Henry C. Jeanette	Reg. No. 30,856
Eric S. Dicker	Reg. No. 31,699	Susan Lee	Reg. No. 30,653
Norman C. Dulak	Reg. No. 31,608	Anita W. Magatti	Reg. No. 29,825
Cynthia L. Foulke	Reg. No. 32,364	John J. Maitner	Reg. No. 25,636
Robert A. Franks	Reg. No. 28,605	Joseph T. Majka	Reg. No. 30,570
James M. Gould	Reg. No. 33,702	Edward H. Mazer	Reg. No. 27,573
Richard J. Grochala	Reg. No. 31,518	James R. Nelson	Reg. No. 27,929
Thomas D. Hoffman	Reg. No. 28,221	Paul A. Thompson	Reg. No. 35,385
Carl W. Battle	Reg. No. 30,731	Arthur Mann	Reg. No. 35,598
Sheela Mohan-Peterson	Reg. No. P-41,201	Jaye P. McLaughlin	Reg. No. P-41,211

Send Correspondence to:

Edwin P. Ching  
DNAX Research Institute  
901 California Avenue  
Palo Alto, CA 94304-1104

Direct Telephone Calls to:

Name: Edwin P. Ching  
Telephone No.: (415) 496-1204  
Facsimile No.: (415) 496-1200

FULL NAME OF 1ST OR SOLE INVENTOR	FAMILY NAME	FIRST GIVEN NAME	SECOND GIVEN NAME
	Wang	Wei	
RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	Palo Alto	CA	People's Republic of China
POST OFFICE ADDRESS	POST OFFICE ADDRESS	CITY	STATE & ZIP CODE/COUNTRY
	3090 Middlefield Road	Palo Alto	CA 94306 / USA

FULL NAME OF 2ND JOINT INVENTOR	FAMILY NAME	FIRST GIVEN NAME	SECOND GIVEN NAME
	Gish	Kurt	C.
RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	San Francisco	CA	USA
POST OFFICE ADDRESS	POST OFFICE ADDRESS	CITY	STATE & ZIP CODE/COUNTRY
	974 Hayes Street	San Francisco	CA 94117 / USA

Wei WANG, et al.; USSN No.: to be assigned;  
Filed: July 3, 1997; Express Mail Label: EM 161 836 205 US

FULL NAME OF 3RD JOINT INVENTOR	FAMILY NAME Schall	FIRST GIVEN NAME Thomas	SECOND GIVEN NAME J.
RESIDENCE & CITIZENSHIP	CITY Menlo Park	STATE OR FOREIGN COUNTRY CA	COUNTRY OF CITIZENSHIP USA
POST OFFICE ADDRESS	POST OFFICE ADDRESS 2050 Mills Avenue	CITY Menlo Park	STATE & ZIP CODE/COUNTRY CA 94025 / USA

FULL NAME OF 4TH JOINT INVENTOR	FAMILY NAME Vicari	FIRST GIVEN NAME Alain	SECOND GIVEN NAME
RESIDENCE & CITIZENSHIP	CITY Mountain View	STATE OR FOREIGN COUNTRY CA	COUNTRY OF CITIZENSHIP France
POST OFFICE ADDRESS	POST OFFICE ADDRESS 2250 Latham Street, #54	CITY Mountain View	STATE & ZIP CODE/COUNTRY CA 94040 / USA

FULL NAME OF 5TH JOINT INVENTOR	FAMILY NAME Zlotnik	FIRST GIVEN NAME Albert	SECOND GIVEN NAME
RESIDENCE & CITIZENSHIP	CITY Palo Alto	STATE OR FOREIGN COUNTRY CA	COUNTRY OF CITIZENSHIP USA
POST OFFICE ADDRESS	POST OFFICE ADDRESS 507 Alger Drive	CITY Palo Alto	STATE & ZIP CODE/COUNTRY CA 94036 / USA

I further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Signature of First Inventor	Signature of Second Inventor	Signature of Third Inventor
Wei Wang	Kurt C. Gish	Thomas J. Schall
Date	Date	Date

Signature of Fourth Inventor	Signature of Fifth Inventor
Alain Vicari	Albert Zlotnik
Date	Date